

## The Kampar pinnacles: Sedimentology and depositional environment

KHOR WEI CHUNG<sup>1,\*</sup>, MOHD SHAFEEA LEMAN<sup>2</sup>, NURUL ‘AMALINA BINTI MD NOR<sup>3</sup>,  
MUHAMMAD ASHAHADI DZULKAFI<sup>2</sup>, MAT NIZA BIN ABDUL RAHMAN<sup>3</sup>,  
MOHD AL-FARID ABRAHAM<sup>1</sup>

<sup>1</sup> Fakulti Sains & Teknologi, Universiti Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu, Sabah, Malaysia

<sup>2</sup> Geology Programme, Department of Earth Sciences and Environment,

Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

<sup>3</sup> Jabatan Mineral dan Geosains Malaysia, Jalan Sultan Azlan Shah, 31400, Ipoh, Malaysia

\* Corresponding author email address: [khorweichung@ums.edu.my](mailto:khorweichung@ums.edu.my)

**Abstract:** The Kinta Valley of Perak, Malaysia is a geologically significant region characterized by Paleozoic carbonate deposits, karst landscapes, and a rich fossil record. This study focuses on the sedimentological description and depositional interpretation of carbonate pinnacles once exposed at Putra Terminal, Kampar, which have since been demolished due to urban development. A total of 26 pinnacles were surveyed, with seven logged in detail to construct two continuous lithostratigraphic sections. Five facies were identified: stylonitic limestone (M), laminated limestone (LL), oval detritus limestone (F), bioclastic limestone (BL), and brecciated limestone (Bc). The facies association indicates a dominantly shallow-marine depositional setting ranging from peritidal to a classic reef environments, with episodic higher-energy influences. Fossil assemblages, including gastropods, horn corals, and possible fusulinids, provide evidence for a Late Carboniferous to Permian age. Diagenetic overprinting, particularly dolomitization, was pervasive and restricts microfacies analysis. These results contribute to the broader understanding of the Kinta Limestone Formation and underscoring the importance of documenting rapidly disappearing geological sites within Malaysia’s karst landscapes.

**Keywords:** Kinta Valley, limestone, sedimentology, pinnacles

### INTRODUCTION

The Kinta Valley, located in Perak, Malaysia, has a rich history of geological usage over the past century, largely driven by its abundant tin deposits. During the late 19<sup>th</sup> and early 20<sup>th</sup> centuries, the valley became one of the world’s largest alluvial tin mining areas, significantly contributing to Malaysia’s economy (Lehmann, 2021). This tin boom spurred the development of infrastructure and construction not only in Ipoh but also in smaller towns such as Batu Gajah and Kampar. Post-independence, as tin prices declined and the industry waned, the region diversified its geological applications. The karst landscape of limestone hills and caves gained prominence, leading to the development of geoheritage sites and geotourism (Adnan Jusoh & Yunus Sauman Sabin, 2019). The study of Kinta Valley’s geology, especially its stratigraphy, is crucial for understanding its geological history and resources potential (Aydin, 2000; Brookfield, 2004),

aiding in mineral exploration, construction projects, and enhancing geotourism (Mohd Shafeea Leman *et al.*, 2007; Nordiana Mohd Muztaza *et al.*, 2025). The Kinta Valley is geologically notable for its extensive karst landscape, characterized by limestone hills and numerous caves formed through the dissolution of carbonate rocks. The limestone here primarily belongs to the Kinta Limestone Formation, which dates to the Paleozoic era, making it over 250 million years old. This formation includes significant deposits of limestone and dolomite while being flanked by younger granite (Lee, 2009).

Notable sites like Gua Tempurong, Sam Poh Tong, Perak Tong and Kek Lok Tong caves attracted tourists, emphasizing conservation and educational aspects of geology. The karstic hills in this valley are characterized by the presence of dolines, caves, and caverns, staged waterflow platform with needle-shaped stalactites hanging from the ceilings and stalagmites rising from

the cave floors (Ros Fatimah & Ibrahim, 2003; Pierson, 2009; Choong *et al.*, 2014). Kinta Valley geological formations have been used in construction and industry, and today, the legacy of tin mining is preserved in museums and heritage trails, while the natural beauty of the limestone formations continues to attract visitors worldwide.

This study aims to give a detailed description and comprehensive sedimentological analysis of the carbonate pinnacles exposed at Putra Terminal, Kampar, Perak. These carbonate pinnacles are a distinctive feature of the local karst landscape, representing ancient reef structures that have been exposed through erosion. Studying these formations provides valuable information about the paleoenvironment and geological history of the region. Moreover, these rocks have been demolished for housing development, making these reported sites not available for future visits.

## OVERVIEW OF KINTA VALLEY

Kinta Valley is located at the Western Belt of Peninsula Malaysia in the state of Perak. It is drained by the Kinta River, which originates from Mount Korbu at Ulu Kinta flowing southward to Tanjung Rambutan, Kampung Gajah and finally join up with the Perak River and at Kampung Bandar.

The geology of Kinta Valley is characterized by a combination of complex stratigraphy primarily consisting of sedimentary and volcanic rock units. It is bounded by Kledang Granite Range north and eastward. While the western margin is bounded by Kati Formation and Bujang Melaka (Figure 1). The dominating lithological units present in Kinta are limestone, shaley limestone, sandstone and mudstone.

The nomenclature of Kinta Limestone Formation is established based on localities such as mines and settlements. Based on Suntharalingam (1968); Rajah (1979) and Henri & Ibrahim Amnan (1995), the stratigraphy of Kinta Limestone comprises shallow water sediments that can be distinguished into six lithological units. These lithological units are i) "Kim Loong Beds No. 1", ii) "Thye On Beds", iii) "Kuan On Beds", iv) "Kim Loong Beds No. 3", v) "Nam Loong Beds", and the youngest, vi) "H.S. Lee Beds" (Figure 2). "Kim Loong Beds" is reported to be cream-coloured dolomite limestone. This unit is assigned to age from Givetian, Lower Devonian to Silurian, in particular Givetian in age (Henri & Ibrahim Amnan, 1995), making it the oldest unit. The thickness is estimated to be 600 m and distributed mainly around Gunung Kanthan. Tabulate corals and conodonts reportedly found in the unit (Alexander & Muller, 1963; Metcalfe, 2000). "Kim Loong Beds" was also reported to be similar with Chemor Limestone by Gobbett (1971). Overlying the "Kim Loong Beds" is the 150 m thick "Thye On Beds". This unit consists of massive grey recrystallized calcitic limestone distributed at the southern end of Kinta Valley. Several fossils were reported, including conodonts, Murchisoniidae gastropods, onocerid nautiloids, ambocellid brachiopod stringocephalus perakenis and tabulate corals (Gobbett, 1966; Metcalfe, 2000). "Kuan On Beds" is aged Lower Carboniferous with estimated thickness of 500 m and is exposed near Batu Gajah, west of Kampar, in the vicinity of Kampung Sungai Kroh and northeast of Malim Nawar. "Kuan On Beds" is predominantly greyish and thinly bedded recrystallized calcite




Figure 1: Geological map of Kinta Valley (adapt from JMG, 2024).

Age	Thick-ness	Member	Lithology	Description	Fossil	Locality
Lower Permian	20 m	H.S. Lee Beds		Biohermal limestone	i. Gastropod, ostracods, cidaroid, corals ii. Chiton valves, bivalves, nautiloid	i. East of Batu Gajah
	20 m			Grey bioclasts & fusulinid limestone	i. Larger foram ii. Broken or complete gastropod shells	i. East of Batu Gajah
	60 m	Nam Loong Beds		Impure carbonaceous brachiopod - polyzoan limestone	i. Brachiopods & gastropods ii. Small brachiopods (Gobett, 1970)	i. East of Batu Gajah
	100 m			Crinoidal	i. Crinoid cups & columnal ii. Small brachiopods	i. West of Kampar
Upper? Carboniferous	100 m	Kim Loong Beds No. 3		Pyritic black shale & argillaceous mudstone	Unfossiliferous	i. West of Kampar
Lower Carboniferous	~500 m	Kuan On Beds		Thin bedded grey recrystallized calcite limestone with dolomite beds interbedded with calcareous and carbonaceous shale	i. Gastropods, bivalve Schizodus (Gobett, 1970) ii. Corals, gastropod (Jones, 1966)	i. Near Batu Gajah ii. West of Kampar near Kampung Sungai Kroh & NE of Malim Nawar
Middle? to Upper Devonian	150 m	Thye On Beds		Massive grey recrystallized calcitic limestone	i. Conodonts (Metcalfe, 2000) ii. Murchisoniidae gastropods, onocerid nautiloids, & amboceliid brachiopod stringocephalus perakenis (Gobett 1966) iii. Tabulate corals (Gobett, 1970)	i. Southern end of Kinta Valley
Lower Devonian to Silurian?	~600 m	Kim Loong Beds No. 1		Cream-coloured dolomite	i. Tabulate corals (Jones, 1973) ii. Conodonts (Alexander & Muller, 1963; Metcalfe, 2000)	i. Gunung Kanthan, or Chemor Limestone

Figure 2: General stratigraphy of Kinta Valley and study area (after Gobbett, 1973).

limestone and ooids limestone with minor interbedded dolomite, calcareous and carbonaceous shale. Gastropods, bivalve Schizodus and corals have been discovered in this unit (Gobbett, 1971). Overlying the “Kuan On Beds” is the 100 m thick Upper Carboniferous aged “Kim Loong Beds No. 3”. This unit consists of unfossiliferous sandstone, pyritic black shale and argillaceous mudstone. The youngest two units, with the latter being younger, “Nam Loong Beds” and “H.S. Lee Beds” are both reported to be Lower Permian in age with relative thickness of less than 100 m. These two units are rich in fossils, namely, gastropod, ostracods, cidaroids, corals, chiton valves, bivalves, nautiloid, larger foram, smaller brachiopod and crinoid (cups and columnal).

The arenite series in Kinta Valley was initially described and named by Ingham & Bradford (1960) as Kati Beds. Later the term Kati Formation was introduced by (Foo, 1983; Alkhali & Sum, 2015). This Carboniferous to Upper Permian aged unit is exposed westward of Batu Gajah. It is reported that Kati Formation is highly weathered and folded. The lithological unit comprises massive sandstone, interbedded sandstone & shale and greyish shale with iron concretion (Khan & Gámez Vintaned, 2022).

**METHOD AND MATERIALS**

For sedimentological and stratigraphical studies, the limestone pinnacles in the area were measured and correlated from a lithostratigraphic perspective. The exposed limestone pinnacles, located at 4.316789°N, 101.130865°E near Terminal Putra Kampar were examined for their sedimentological characteristics and fossil content. A total of 26 pinnacles were mapped, and seven correlatable pinnacles were logged to produce





Figure 3: Overview of the exposed pinnacles. The logged seven sections identified (P1 – P7).

continuous sections (Figure 3 & 4). The fundamental elements and features of carbonate facies, facies model and their generally accepted interpretations, were adapted from Tucker (1985) to support further interpretation. This study attempts to characterize the carbonate through physical characteristics (Flügel, 2004; Tucker, 2009). In these studies, three shallow-marine carbonate settings that may evolve into one another were proposed: 1) platforms, 2) shelves, and 3) ramps.

The dataset used in this research can be obtained from Khor *et al.* (2025). It contains recorded sedimentary logs, XRD results, petrography data, photographs of pinnacles, lithologies and fossils from the study area.

**RESULT AND DISCUSSIONS**

The exposed rocks are rather monotonous, relatively crystallized and thinly bedded succession. In this paper, the facies is classified based the physical observation and sedimentary characteristics. Likewise, facies classification based on microphotographs/microfacies were done by



**Figure 4:** Two sections of continuous logs (13.5 m and 2 m thick). Section 1 consist of pinnacles P4, P5, P6 and P7. Section 2 consist of pinnacles P2 and P3.

**Table 1:** List of the five facies identified in study area.

No	Facies	Code	Color	Texture	Description	Fossils	Boundaries	Thickness
1	Stylolitic Limestone Facies	M	Greyish white and minor yellowish hue	Mudstone	Structureless, stylolitic	Gastropod	Irregular and planar lower contacts	20 cm to 4 m
2	Bioclastic Limestone Facies	BL	Grey micrite and greyish white bioclasts	Wadestone and minor grainstone	Mud-supported, no imbrication	Tabulate coral, isolated coral, crinoid stem?	Irregular/ scouring bottom contact with LL facies	1 m in Pinnacle 4
3	Oval Detritus Facies	F	Grey micrite and greyish white bioclasts	Wadestone	Imbricated, rounded clasts include 1-1.5 cm oval detritus with minor rounded clasts – averaging 1 cm in diameter	Tabulate coral, fusulinid, isolated coral	Planar lower contact with M	60 cm thick in Pinnacle 2
4	Brecciated Limestone Facies	Bc	Greyish white and greyish brown	Breccia	No imbrication, polymictic, angular clasts less than 5 cm in diameter, matrix supported, clasts include both laminated and structureless limestone	Wood fragment 10 cm long	Planar top contact with Bc facies	Wedge-like, 40 cm thick in Pinnacle 4
5	Laminated Limestone Facies	LL	Darker grey micrite/fine materials with white calcite bands. Lamina-tions are dark grey	Mudstone	Lamination trough stratification	-	Sharp contact and transition with M facies	Averages 30 cm in thickness

Poon *et al.* (2022). Five facies identified are i) Stylolitic limestone facies (M), ii) Bioclastic limestone facies (BL), iii) Oval detritus limestone facies (F), iv) Brecciated limestone facies (Bc), and v) Laminated limestone facies (LL) (Figure 5, Table 1).

### Stylolitic limestone facies (M)


This facies is composed of structureless stylolitic limestone, predominantly greyish-white in color. The outer portions of the rock often exhibit a yellowish hue due to weathering. Bed thickness varies from 20 cm to 4 m, with some beds appearing graded to the naked eye. Hand samples reveal rhombic cleavage. The stylolite sutures are seam- or horsetail-like with amplitude not exceeding 1 cm and commonly infilled with greyish mudstone. Rounded clasts are occasionally observed at the base of the beds, while scoured and planar lower contacts are common. The upper contact is typically gradational into the laminated limestone facies (LL). Facies M is commonly found in most pinnacles. Gastropod fossils are present within this facies. Stylolitic structures in limestone represent pressure dissolution that occur due to overburden pressure resulting in bed thickness, permeability and porosity reduction (Norman, 2015; Magni *et al.*, 2025). Seam-like stylolites are interpreted to be from an impermeable layer (Sadd & Alsharhan, 2000; Gomez-Rivas *et al.*, 2022) along with scoured bottom contacts indicate the participation of wave action. The presence of gastropod which are more diverse in shallower environment, possibly mid-shelf to intertidal, eliminate the likelihood of open-ocean (Dahl, 2012).

### Bioclastic limestone facies (BL)

This facies is composed of bioclasts which include rugose corals, isolated corals, and possibly crinoid stems. According to Dunham's classification, the rock is primarily mud-supported which can be classified as a wackestone. The facies is one meter thick and occurs exclusively in Pinnacle 4 (P4) (Figure 4). In some instances, certain sections contain a higher percentage of grains, locally transitioning into grainstone. This facies portrays similarities to facies F, except for the presence of imbrication. This facies shows similarity to the described Nam Long Beds in Gobbett (1973) which is rich with crinoid, foraminifera, brachiopods, gastropods and fusulinids. Based on Henri & Ibrahim Amnan (1995) this facies is deposited in calm environments and are habitual to such organisms.

### Oval detritus facies (F)

In Pinnacle 2 (P2), this facies reaches a thickness of 60 cm, forming wedge-like bed. It is dominated by elongated, wheat-like oval detritus, averaging 1-1.5 cm in length, with some exceeding 2 cm, set within micrite matrix. Minor rounded clasts, averaging 1 cm in diameter, are also present. The primary distinction between this facies and the bioclastic limestone facies (BL) lies in its detrital composition and structure. The oval detritus, possibly *Triticites*, representing recrystallized medium to large fusulinid fossils, which were widespread and peaked during Carboniferous (Sims, 2020; Huang *et al.*, 2023). Vertical line patterns (fusiform shape) observed in some specimens may



**Figure 5:** Photos of identified facies. A & B) laminated with platy calcite bands (LL), C) bioclastics limestone with rounded clasts, D & E) laminated greyish limestone, and F) horse-tailed stylolite.

correspond to chamber accretion, as indicated by groove features known as furrows. Additionally, elongated spicules within the facies could represent isolated horn corals, which were abundant before the K-T extinction. The occurrence of this facies as a single wedge-like bed, along with the random orientation of detrital grains, further supports evidence of reworking before final deposition. However, larger fusulinids are known to delve in high energy environments, particularly wave-dominated zones and the random orientation could be a common occurrence in such environments (Sarwary *et al.*, 2022).

### Brecciated limestone facies (Bc)


This facies consists of broken, angular carbonate clasts (Figure 7D), with rare instances of clast-supported texture within a predominantly matrix-supported framework, suggesting *in-situ* deposition with minor fine sediment influx. It is classified as a polymictic, matrix-supported breccia, with observed clasts including laminated limestone and greyish structureless fragments. A 10 cm long wood fragment imprint is also present. Diagenetic alterations are evident, with some minor clasts appearing more rounded and exhibiting cul-de-sac pores. The brecciated clasts range from millimeters to 4 cm in diameter and remain predominantly angular. It can form due to failed cave, *in-situ* formation or lag deposit (Scholle *et al.*, 1992).

### Laminated limestone facies (LL)

This facies consists of light greyish limestone, averaging 30 cm in thickness, with laminations of darker bands (Figure 6). It is composed predominantly of fine-grained carbonate mud with minor platy bedding and is classified as a wackestone. The laminations are not sharply defined and occasionally appear trough-shaped, with diameters ranging from 2-5 cm and heights of up to 20 cm, suggesting minor wave influence in a low-energy environment. A gradual and transitional contact with the stylolitic limestone facies (M) is commonly observed. This concaved trough is interpreted as bioturbation by trace fossils in a shallower water environment, possibly lagoonal area (Kamal Roslan Mohamed & Che Aziz Ali, 2010; Alonso-Zarza *et al.*, 2011). The laminated fine materials in this facies is likely consisting of algae, organic materials, carbonate mud or possibly siliceous mud. The lamination observed indicates low-energy and rapid, alternating input suggesting deposition in a protected lagoonal or tidal flat environment.

## FOSSILS

The fossils identified and available in this dataset are i) isolated horn coral, ii) gastropod, iii) fragmented tabulate coral, possibly iv) fusulinid, and v) crinoid stem (Figure 7).



**Figure 6:** Photos of logged pinnacles. A) Pinnacle P4 – Faulted pinnacle with bioclastic facies overlying stylolitic and laminated limestone facies at the right, B) Fractured pinnacle with orthogonal joints C) Pinnacle P6 – stylolitic facies overlying laminated limestone facies, and D) Pinnacle P7 – laminated limestone facies.

#### i. Isolated horn coral


The other possibility is that it can be sponge skeletal spicules. However, the fact that the vertical lines feature on the fossils and larger size rule out the possibility of being megascleres sponge skeletal spicules (cf. Łukowiak *et al.*, 2022). Likewise, preserved bryozoans are usually the y-shape brunch, fan- and net-like cast (cf. Ernst *et al.*, 2024).

#### ii. Fusulinid

The identification of fusulinid in facies F is lacklustre as it lacks petrography evidence. The early assumption obtained from this study is that the oval detritus in facies F is most possibly fusulinids. Fusulinid is widely spread especially in Permian aged limestone. Fusulinid is also reported in the Permian aged limestone such as Kubang Pasu Formation (Shirin Fassihi *et al.*, 2024) and Triassic aged limestone in Gua Musang Formation.

## DIAGENESIS

Dolomitization in Kinta Valley has been well described by (Poon *et al.*, 2022) from metamorphosed dolomite, early and late dolomitization to sucrosic dolomite. The extensive



**Figure 7:** A) gastropod, B) isolated coral, horn? C) wackestone, tabulate coral clast in, D) *in-situ?* breccia, E) Elongated fusiform and rounded detritus wackestone (fusulinid and corals?), F) wackestone - redeposited bioclasts with possible disarticulated brachiopods, carbonaceous fragments.

recrystallization, dolomitization and occurrence of host calcareous mudstone (HCM) were said to be the result of thermal events (Henri & Ibrahim Amnan, 1995). Poon *et al.* (2022) further noted that multiple phases of dolomitization were caused by deep-circulating crustal hydrothermal fluids associated with tectono-thermal events. Occurrence of Sr element based on the XRD done on stylolitic limestone facies (M) suggest diagenesis through fluid development and selective dolomitization (Nurul Afifah Mohammad Zahir *et al.*, 2020). The carbonate pinnacles studied are dolomitic and recrystallized. Thin section observation identified dormant partial or selective dolomitization. It is interpreted that episodic dolomitization in Kinta Valley took place during Permian to Cenozoic (Ramkumar *et al.*, 2019). During these periods the paleoenvironment transitioned from marine to terrestrial. The presence of Sr plays an important role in dolomitization (Veizer, 1978). Higher value of Sr could indicate the presence of aragonite during dolomitization process than those originating from calcite.

### DEPOSITION ENVIRONMENT

The discussion is based solely on the physical characteristics observed at the pinnacles. Microphotography is not possible with the degree of crystallization present in the prepared thin sections. The facies identified in this study show similarities to the Lower Permian aged Nam Loong and H.S. Lee Beds as compiled in Gobbett (1971). Previous studies suggest that the Kinta Limestone was deposited in a range of marine environments, from deep-marine lower slope settings to shallower facies such as platform

margins and lagonal areas toward the east (Pierson *et al.*, 2011). Gebretsadik *et al.* (2015) interpreted parts of the Kinta Valley as representing slope and intrabasin-margin settings, based on the presence of chert, the distribution of conodonts in the northern sector, and southward thinning of sedimentary beds. It is also described that Kinta Valley limestone was deposited in a pelagic to hemipelagic environment. Henri & Ibrahim Amnan (1995) interpreted a calm and subtidal environment for limestones around Kampar region. More recent works, including Gebretsadik *et al.* (2014) and Mahendran (2021) suggested rimmed platform model for the limestone in Kampar area.

The pinnacles in this study show the dominance of stylolitic limestone (M) and laminated limestone (LL) with subordinate oval detritus limestone (F) and underlain by bioclastic limestone (BL). Brecciated limestone (Bc) should not be taken into consideration as it forms independently from localized process such as collapsed cave, failed bank or could potentially form at the upper slope (Figure 8). The facies assemblage documented indicates deposition on a classic reef carbonate model, positioned along the western margin of the Kinta Valley during the Late Carboniferous-Permian. The succession is dominated by low-energy limestones (stylolitic limestone, M facies; laminated limestone, LL) interbedded with localized bioclastic and fusulinid-rich horizons (Bc and F), suggesting a reef interior setting punctuated by episodic high energy events. Although facies LL indicates low-energy settings, its laminations can form in various environments, including restricted lagoon, tidal flats or even deep basins under anoxic conditions. Facies F records




Figure 8: Proposed classic reef model carbonate model for Kampar Limestone. Modified from Wilson (1975).

possible fusulinid fossil content where it indicates a wide range of depositional environment, occurring in shallow water to deeper areas of a carbonate platform setting (Avendaño-Pazos *et al.*, 2024).

Together, these facies suit a classic reef system (Figure 8), where energy is concentrated along the reef, rapidly decreasing landward into a protected lagoon. The absence of deepening-facies, storm-dominated outer ramp deposits, siliciclastic facies or gradual proximal-distal transitions further argues against a unrimmed platform model.

### CONCLUSION

This study aims to preserve, document, and share valuable geological information on the now-demolished limestone pinnacles, which were removed due to ongoing urban development. The findings are significant as they provide one of the few remaining sedimentological records of the Kampar carbonate pinnacles, contributing essential data for reconstructing the depositional history and paleoenvironments of the Kinta Limestone.

The rock sequence recorded in this study suggests a repetitive occurrence of stylolitic limestone (facies M) and laminated limestone (facies LL), interpreted to have formed in wide range of settings including peritidal, lagoonal to tidal flat. The subordinate occurrence of oval detritus limestone (facies F) and bioclastic limestone (facies Bc) in isolated pockets indicates possible reworking or localized periods of biogenic accumulation.

However, further evidence is required, as stylolitic limestone facies can develop as a result of diagenesis in a wide range of depositional settings. Additional studies focusing on the types of stylolites present may provide better

insight into the original characteristics of limestone prior to diagenesis. It is likely that formation of stylolites occurred between two units with contrasting porosity. Likewise, laminated limestone facies also deposit in a wide range of environments. Distinguishing between ramp, platform and shelf would require longer vertical sections with clear bathymetric gradient indicators, microfacies and biofacies zonation. A classic reef model is proposed for these pinnacles.

From a rock fabric perspective, more extensive studies on the fossil content and its incorporation into facies characterisation could help overcome observational challenges posed by intense recrystallization within these limestones.

### ACKNOWLEDGEMENT

The authors would like to thank the reviewers for their valuable comments. Sincere thanks are also extended to UKM Geology for preparing the thin sections and to JMG (Jabatan Mineral dan Geosains Malaysia) for organizing the field excursion in February 2024.

### AUTHORS CONTRIBUTION

KWC - Drafting and data acquisition; MSL - data acquisition, drafting and thin sections; NABMN - field organizer and mapping; MAD - data acquisition, paleontological studies and thin sections; MNBAR - field organizer and data acquisition; MAA - XRD and geochemistry.

### ETHICS

This study does not involve humans or animals as test subjects. Data collection and field excursions were

conducted in accordance with government and Shariah laws.

### CONFLICT OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### REFERENCES

- Adnan Jusoh & Yunus Sauman Sabin, 2019. Analysis of Local Tourists' Level of Knowledge on Archaeotourism Sector in Kinta Valley, Perak (Malaysia). *International Journal of Innovation, Creativity and Change*, 6(2), 116 – 131.
- Alexander, J.B. & Muller, K.J., 1963. Devonian Conodonts in Stratigraphic Succession of Malaya. *Nature*, 197, 681.
- Alkhali, H.A. & Sum, C.W., 2015. The Kati Formation: A Review. *Proceedings of the International Conference on Integrated Petroleum Engineering and Geosciences (ICIPEG 2014)*, 303-312.
- Alonso-Zarza, A.M., Genise, J.F. & Verde, M., 2011. Sedimentology, diagenesis and ichnology of Cretaceous and Palaeogene calcrites and palustrine carbonates from Uruguay. *Sedimentary Geology*, 236, 45-61.
- Avendaño-Pazos, J.J., Torres-Martínez, M.A., Lara-Peña, R.A. & Navas-Parejo, P., 2024. Biostratigraphy of fusulinids from the lower Permian of central Sonora, Mexico. Paleoenvironmental implications. *Journal of South American Earth Sciences*, 148, 105133.
- Aydin, A., 2000. Fractures, faults, and hydrocarbon entrapment, migration and flow. *Marine and Petroleum Geology*, 17, 797-814.
- Brookfield, M., 2004. *Principles of Stratigraphy*. Blackwell Publishing, Oxford. 352 p. ISBN 13: 9781405111645.
- Choong, C.M., Sautter, B., Pubellier, M., Menier, D., Chow, W.S. & Askury Abd Kadir, 2014. Geological features of the Kinta Valley. *Platform*, 10(2), 13 p.
- Dahl, R.M., 2012. A Paleoenvironmental Analysis of Gastropod from the Middle Ordovician, Ibex Region, Utah. Unpublished thesis, University of California Riverside.
- Ernst, A., Seroby, V. & Danelian, T., 2024. Biostratigraphic, palaeoenvironmental and palaeobiogeographic implications of bryozoan fauna from the Upper Devonian sequences of Armenia. *Geobios*, 85(1777-5728), 10–18. <https://doi.org/10.1016/j.geobios.2024.02.001>.
- Flügel, E., 2004. *Microfacies of Carbonate Rocks: Analysis, Interpretation and Application*. Springer, Berlin, Heidelberg, New York. 976 p.
- Foo, K.Y., 1983. The Paleozoic Sedimentary Rocks of Peninsular Malaysia. Workshop on Stratigraphic Correlation of Thailand and Malaysia. Haad Yai, Thailand, 8-10 Sep. 1983.
- Gebretsadik, H.T., Chow, W.S. & Hunter, A.W., 2015. Discovery of Upper Devonian-Lower Carboniferous conodonts in the Kinta Limestone, Western Belt of Peninsular Malaysia: Implication for continuous sedimentation in the Paleotethys. *Proceedings of the International Conference on Integrated Petroleum Engineering and Geosciences*, 291-312.
- Gebretsadik, H.T., Hunter, A.W. & Chow, W.S., 2014. Depositional Environment of the Kinta Limestone, Western Peninsular Malaysia. *Proceedings of the AAPG International Conferences and Exhibition* 36.
- Gobbett, D.J., 1966. The brachiopod genus *Strigocephalus* from Malaya. *Journal of Paleontology*, 40, 1345-1348.
- Gobbett, D.J., 1971. Joint patterns and faulting in Kinta, West Malaysia. *Bulletin of the Geological Society of Malaysia*, 4, 39–48. <https://doi.org/10.7186/bgsm04197103>.
- Gobbett, D.J., 1973. Geology of the Malay Peninsula; (West Malaysia and Singapore). In: C.S. Hutchinson (Ed.). Retrieved September 9, 2025, from Internet Archive website: <https://archive.org/details/geologyofmalaype0000gobb>.
- Gomez-Rivas, E., Martin-Martin, J.D., Bons, P.D., Koehn, D., Griera, A., Trave, A., Llorens, M.G., Humphrey, E., & Neilson, J., 2022. Stylolites and stylolite networks as primary controls on the geometry and distribution of carbonate diagenetic alterations. *Marine and Petroleum Geology*, 136, 1 – 14. <https://doi.org/10.1016/j.marpetgeo.2021.105444>.
- Henri, F. & Ibrahim Amnan, 1995. Biostratigraphy of the Kinta Valley, Perak. *Bulletin of the Geological Society of Malaysia*, 38, 159-170.
- Huang, H., Jin, X., Shi, Y., Zheng, J., & Yan, Z., 2023. New fusulinid assemblages from the Changning-Menglian Belt in western Yunnan, China and their paleogeographic implications. *Journal of Asian Earth Sciences*, 256, 105807. <https://doi.org/10.1016/j.jseas.2023.105807>.
- Ingham, F.T. & Bradford, E.F., 1960. The Geology and Mineral Resources of the Kinta Valley, Perak. *Federation of Malaya Geological Survey, Ipoh*. 347 p.
- JMG, 2024. Geological Map and Lineament. *Jabatan Mineral dan Geosains (Malaysia)*.
- Kamal Roslan Mohamed & Che Aziz Ali, 2010. Geologi dan landscape Bukit Batu Putih, Taman Rimba Kenong, Pahang. *Bulletin of the Geological Society of Malaysia*, 56, 95-105. <https://doi.org/10.7186/bgsm56201014>.
- Khan, M.E.N. & Gámez Vintaned, J.A., 2022. Boundaries and lithofacies of the Kati Formation (SW Perak, Peninsular Malaysia). *IOP Conf. Ser.: Earth Environ. Sci.* 1003. <https://doi.org/10.1088/1755-1315/1003/1/012020>.
- Khor, W.C., Mohd Shafiea Leman, Mohamad Ashahadi Dzulkafli, Nurul' Amalina binti Md Nor & Mat Niza Abdul Rahman, 2025. Carboniferous – Permain Kinta Limestone Pinnacles, Kampar, Perak, Malaysia. *Mendeley Data*, V1. <https://doi.org/10.17632/rvdcjsj78j8.1>.
- Lee, C.P., 2009. Paleozoic Stratigraphy. In: Hutchison, C.S. & Tan, D.N.K. (Eds.), *Geology of Peninsular Malaysia*. Geological Society of Malaysia, Kuala Lumpur. 479 p.
- Lehmann, B., 2021. Formation of tin ore deposits: A reassessment. *Lithos*, 402-403, 1-4. <https://doi.org/10.1016/j.lithos.2020.105756>.
- Lukowiak, M., Soest, V.R., Klautau, M., Pérez, T., Pisera, A., & Tabachnick, K., 2022. The terminology of sponge spicules. *Journal of Morphology*, 283(12), 1517–1545. <https://doi.org/10.1002/jmor.21520>.
- Magni, S., Martin-Martin, D.J., Bons, P.D., & Gomez-Rivas, E., 2025. Stylolites in Carbonate Rocks: Morphological Variability According to the Host Rock Texture. *Minerals*, 15(132), 1 – 23. <https://doi.org/10.3390/min15020132>.
- Mahendran, S., 2021. Geologi Batu Kapur Kampar, Perak. Unpublished thesis, Universiti Kebangsaan Malaysia.
- Metcalfe, I., 2000. The Bentong-Raub Suture Zone. *Journal of Asian Earth Sciences*, 18, 691-712. <https://doi.org/10.1016/>

S1367-9120(00)00043-2.

- Mohd Shafeea Leman, Ibrahim Komoo, Kamal Roslan Mohamed, Che Aziz Ali & Tanot, U., 2007. Geopark as an answer to geoheritage conservation in Malaysia, the Langkawi Geopark case study. *Bulletin of the Geological Society of Malaysia*, 53, 95-102.
- Nordiana Mohd Muztaza, Gopalan, T., Nur Amanina Mazlan, Jamaludin Othman, Nazrin Rahman, Najmiah Rosli, Rosli Saad, Farid Najmi Rosli, Athirah Rosli & Yasir Bashir, 2025. Utilizing Gravity Surveys for Subsurface Feature Identification in Foundation Planning. *Sains Malaysiana*, 54(6), 1465 – 1475.
- Norman, K., 2015. Stylolitization of limestone – A study about the morphology of stylolites and its impacts of porosity and permeability in limestone. Uppsala Universitet. Unpublished thesis.
- Nurul Afiah Mohammad Zahir, Mirza Arshad Beg & Askury Abd Kadir, 2020. Hydrothermal dolomitization on Devonian to carboniferous carbonates in Kinta Valley, Perak, Malaysia: a petrographic study. *Indonesian Journal on Geoscience*, 7(1), 25–39. <https://doi.org/10.17014/ijog.7.1.25-39>.
- Pierson, B., 2009. The limestone hills of the Kinta Valley: a part of Malaysia's geological heritage worth preserving. *First Break*, 27(11), 97-100.
- Pierson, B., Kassa, S., Gebretsadik, H., Askury Abd Kadir, Chow, W.S., Hunter, A. & Harith, Z., 2011. Sedimentology of the Palaeozoic Limestone of the Kinta Valley, Malaysia. *First EAGE South-East Asia Regional Geology Workshop - Workshop on Palaeozoic Limestones of South-East Asia and South China*, December 2011.
- Poon X.H., Nagarajan, R., Ramkumar, M., Ng, T.F., Taib, N.I., Mathew, M.J., Sautter, B., Siddiqui, N.A., & Poppelreiter, M.C., 2022. Geochemical evolution of structure-bedding controlled hydrothermal dolomites of the Kinta Valley, Western Malaysia. *Carbonates and Evaporites*, 37(4). <https://doi.org/10.1007/s13146-022-00802-4>.
- Rajah, S.S., 1979. The Kinta Tinfield, Malaysia. *Bulletin of the Geological Society of Malaysia*, 11, 111-136.
- Ramkumar, M., Siddiqui, N.A., Mathew, M., Sautter, B., Poon, X.H., Nagarajan, R., Breitfeld, H.T., Santosh, M., Menier, D. & Poppelreiter, M., 2019. Structural controls on polyphase hydrothermal dolomitization in the Kinta Valley, Malaysia: Paragenesis and regional tectono-magmatism. *Journal of Asian Earth Science*, 174, 364-380. <https://doi.org/10.1016/j.seaes.2019.02.004>.
- Ros Fatimah, B.H.M. & Ibrahim, K., 2003. The Kinta Valley karst landscape - a national heritage to be preserved. *Bulletin of the Geological Society of Malaysia*, 46, 447-453. <https://doi.org/10.7186/bgsm46200373>.
- Sadd, J. & Alsharhan, A., 2000. Stylolites in lower cretaceous carbonate reservoir, U.A.E. *SEPM Special publication*, No. 69, 185-207. <https://doi.org/10.2110/pec.00.69.0185>.
- Sarwary, M.N., Chaman Shah Alamy, Rahimi, M.L., & Kumar, P., 2022. Interpretation of Depositional Environment of Fusulinid Bearing Middle Permian Succession of Bolula and Khaja Ghar Formation, Bamian Zone, Central Afghanistan. *International Journal of Geosciences*, 13(07), 499–530. <https://doi.org/10.4236/ijg.2022.137027>.
- Scholle, P.A., Ulmer, D.A. & Melim, L.A., 1992. Late-stage calcites in Permian Capitan Formation and its equivalents, Delaware Basin margin, West Texas and New Mexico: evidence for replacement of precursor evaporites. *Sedimentology*, 39, 207-234.
- Shirin Fassihi, Vachard, D., Heinz, P., Meor Hakif Amir Hassan & Azrin Azmi, 2024. A new fusulinid species from the early Permian of northwestern Peninsular Malaysia: Evidence of the Artinskian warming event in the eastern Peri-Gondwanan Sibumasu Block. *Marine Micropaleontology*, 191, 1-15. <https://doi.org/10.1016/j.marmico.2024.102391>.
- Sims, E.R., 2020. Morphometric Analysis of Biostratigraphically Significant Fusulinid Foraminiferan Genus *Triticites* Across the Pennsylvanian-Permian Boundary in the Central and Southwestern United States. Unpublished thesis, Texas A&M University.
- Suntharalingam, T., 1968. Upper Palaeozoic Stratigraphy of the Area West of Kampar, Perak. *Geological Society of Malaysia Bulletin* 1, 1-15.
- Tucker, M.E., 1985. Shallow marine carbonate facies. *Geological Society of London Special Publications*, 18, 147-169.
- Tucker, M.E. 2009. *Sedimentary Petrology: An Introduction to the Origin of Sedimentary Rocks*. Wiley & Sons, New York. 272 p.
- Veizer, J., 1978. Secular variations in the composition of sedimentary carbonate rocks, II. Fe, Mn, Ca, Mg, Si and minor constituents. *Precambrian Res.*, 6(3–4), 381–413. [https://doi.org/10.1016/0301-9268\(78\)90024-4](https://doi.org/10.1016/0301-9268(78)90024-4).
- Wilson, J.L., 1975. *Carbonate Facies in Geologic History*. Springer Verlag, New York. 471 p. <https://doi.org/10.1007/978-1-4612-6383-8>.

*Manuscript received 11 September 2025;*  
*Received in revised form 4 November 2025;*  
*Accepted 21 January 2026*  
*Available online 30 April 2026*