Nature and demise of the Proto-South China Sea

ROBERT HALL* & H. TIM BREITFELD

SE Asia Research Group, Department of Earth Sciences, Royal Holloway University of London, Egham, Surrey, TW20 0EX, United Kingdom *Corresponding author: robert.hall@rhul.ac.uk

Abstract: The term Proto-South China Sea has been used in a number of different ways. It was originally introduced to describe oceanic crust that formerly occupied the region north of Borneo where the modern South China Sea is situated. This oceanic crust was inferred to have been Mesozoic, and to have been eliminated by subduction beneath Borneo. Subduction was interpreted to have begun in Early Cenozoic and terminated in the Miocene. Subsequently the term was also used for inferred oceanic crust, now disappeared, of quite different age, notably that interpreted to have been subducted during the Late Cretaceous below Sarawak. More recently, some authors have considered that southeast-directed subduction continued until much later in the Neogene than originally proposed, based on the supposition that the NW Borneo Trough and Palawan Trough are, or were recently, sites of subduction. Others have challenged the existence of the Proto-South China Sea completely, or suggested it was much smaller than envisaged when the term was introduced. We review the different usage of the term and the evidence for subduction, particularly under Sabah. We suggest that the term Proto-South China Sea should be used only for the slab subducted beneath Sabah and Cagayan between the Eocene and Early Miocene. Oceanic crust subducted during earlier episodes of subduction in other areas should be named differently and we use the term Paleo-Pacific Ocean for lithosphere subducted under Borneo in the Cretaceous. There is good evidence for subduction between the Eocene and Early Miocene below Sabah, and the western limit of Proto-South China Sea subduction was the West Baram Line. The subducted slab can be imaged in the lower mantle using P-wave tomography. There was no subduction beneath Sarawak, SW of the West Baram Line, between the Eocene and Early Miocene where there was terrestrial to marginal marine deposition. The present-day NW Borneo Trough and Palawan Trough are not subduction trenches and these relatively shallow features have different origins. The NW Borneo Trough is largely a flexural response to gravity-driven deformation of the Neogene sediment wedge NW of Sabah whereas the Palawan Trough is the continent-ocean transition at the SE edge of the modern South China Sea.

Keywords: Subduction, Borneo, Sabah, Sarawak, tomography

INTRODUCTION

The late 1960s and early 1970s saw a major change in global tectonic concepts as plate tectonics became the established paradigm. For northern Borneo this was encapsulated by the abandonment of the terminology of undations and geosynclines (e.g. van Bemmelen, 1949; Haile, 1968) and the recognition that subduction had played an important part in the evolution of the region (e.g. Hamilton, 1973; Haile, 1973). However, the links between subduction and the surrounding ocean basins of the South China, Sulu and Celebes Seas (Figure 1) were controversial (see for example, Karig, 1971; Hamilton, 1973; Ben-Avraham & Uyeda, 1973). A clearer picture of Borneo tectonics was slow to emerge because key pieces of information



0126-6187 / © 2017 Geological Society of Malaysia. All rights reserved.



were missing, for example, supposed melanges in Borneo interpreted by Hamilton (1973) were poorly dated, the age of the South China Sea was unknown and was suggested to be Cretaceous by Ben-Avraham & Uyeda (1973) in an early review of geophysical data, and consequently there were quite different ideas concerning the age, location and polarity of subduction zones recorded in Borneo geology.

For the South China Sea this situation had changed by 1980 when Taylor and Hayes identified spreading to have been active from the mid-Oligocene to Early Miocene based on marine geophysical data, principally new magnetic anomaly data, which they followed by reconstructions of the region (Taylor & Hayes, 1983). Their interpretation has subsequently been modified, but largely supported, by later studies (e.g. Briais *et al.*, 1993; Barckhausen & Roeser, 2004; Barckhausen *et al.*, 2014). Ocean drilling (Li *et al.*, 2014, 2015) has subsequently confirmed the age of the oceanic crust of the South China Sea, although the exact ages of the beginning and ending of spreading remain controversial (compare Barckhausen *et al.*, 2014, 2015 with Chang *et al.*, 2015).

Proto-South China Sea

The term Proto-South China Sea (Figure 2) was first used by Hinz et al. (1991) for a large oceanic embayment (Figure 2C) situated in the latest Cretaceous between Indochina and South China in the north and Borneo in the south, that predated the formation of the present-day South China Sea. The oceanic crust of this basin was suggested by Hinz et al. (1991) to have been eliminated by subduction beneath a Borneo microcontinental plate by the Middle Miocene and as subduction proceeded, extension of the continental margin to the north of the Proto-South China Sea led to formation of the modern South China Sea. It is clear that the concept of the Proto-South China Sea in Hinz et al. (1991) was derived from earlier studies. Taylor & Hayes (1980) inferred that a basin older than the South China Sea had existed south of Reed Bank before it was eliminated by subduction beneath Borneo and Taylor & Hayes (1983) showed this as Mesozoic oceanic crust on their Paleocene reconstruction (Figure 2B). Hinz et al. (1991) presented a latest Cretaceous reconstruction (Figure 2C), which is almost identical to the Taylor & Hayes (1983) Paleocene reconstruction (Figure 2B). Holloway (1981, 1982) presented a very similar reconstruction for the Late Cretaceous and Paleocene (Figure 2A).

Paleo-South China Sea

Gatinsky & Hutchison (1986) used the term Paleo-South China Sea for ocean crust north of Borneo in the Jurassic–Cretaceous (150-120 Ma) representing the western part of a Paleo-Pacific ocean. They interpreted it to have been subducted beneath Borneo in the Paleogene (60-40 Ma) as the South China Sea opened, based on Taylor & Hayes (1983). Although the ages adopted were about 20 million years older than interpreted by Taylor & Hayes (1983) this Paleo-South China Sea was a Mesozoic ocean subducted beneath Borneo in the Cenozoic, as confirmed by Hutchison (1989), and therefore similar to the Proto-South China Sea of Hinz *et al.* (1991). Subsequently, the term Paleo-South China Sea has been almost completely abandoned. For example, in later studies Hutchison (1996) used Proto-South China Sea instead. Clift *et al.* (2008) used it for lithosphere subducted beneath Borneo between the Eocene and Early Miocene which all other authors call the Proto-South China Sea.

Features of the Proto-South China Sea

Since the term Proto-South China Sea was introduced by Hinz et al. (1991) it has been used in a broadly similar way by many authors (e.g. Rangin et al., 1990a, 1990b; Rangin & Silver, 1991; Lee & Lawver, 1994, 1995; Tongkul, 1994; Hall, 1996, 2002; Cullen, 2010, 2014), although there have been differences of opinion concerning the size of the ocean basin. Key features of this conceptual Proto-South China Sea are that it formerly occupied the area south of Indochina and South China where the modern South China Sea is situated, its oceanic crust had a Mesozoic age, and it has been completely eliminated by subduction. Subduction began in the Eocene on the southern side of the Proto-South China Sea and ceased in the Early Miocene and the oceanic lithosphere was subducted beneath northern Borneo (Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz et al., 1991), the Cagavan arc (Hinz et al., 1991) and possibly parts of the Philippines (Holloway, 1981, 1982). The subduction zone was shown as extending as far west as offshore Sarawak. The Proto-South China Sea narrowed towards the west and its western end met the Sunda Shelf in a zone of dextral strike-slip faulting. The southward subduction was terminated by collision between continental crust of South China origin and a volcanic arc from Sabah to the Cagayan Ridge. All the models (Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz et al., 1991) showed large counter-clockwise rotation of Borneo during subduction.

However, subsequently the term has also been used for inferred oceanic crust, now disappeared, of quite different age. Some authors have considered that southeast-directed subduction continued until much later than originally proposed, usually based on the misconception that the NW Borneo Trough and Palawan Trough are, or were recently, sites of subduction, a suggestion first made by Hamilton (1973). Hamilton (1979) identified the presentday offshore linear NE-striking troughs as inactive, but relatively young, subduction trenches. He suggested that what he called the NW Borneo Trench, where he interpreted subducted oceanic crust to be visible on seismic reflection profiles, would be interpreted as active were it not for the absence of seismicity, and explicitly labelled his Palawan Trench as "inactivated in Late Miocene". This continues to cause confusion (see Hutchison, 2010b) and has led some authors to consider that subduction beneath northern Borneo continues to the present day (e.g. Tongkul, 1991; Rangin et al., 1999a; Simons et al., 2007). Finally, some authors have challenged the existence of the Proto-South China Sea





Bulletin of the Geological Society of Malaysia, Volume 63, June 2017

of Hinz *et al.* in which the term Proto-South China Sea was first used is almost identical to that of Taylor &Hayes except for the addition of the Borneo microcontinental plate in the SE. 63





Figure 4: Simplified stratigraphy of northern Borneo based on Kirk (1957), Haile (1957), Liechti *et al.* (1960), Wolfenden (1960), Adams & Haak (1962), Haile (1962), Wilson & Wong (1964), Adams (1965), Collenette (1965), Tan (1979), BRGM (1982); Hutchison (1996, 2005, 2010a), Moss (1998), van Hattum *et al.* (2013), Thomson (2013) and Breitfeld (2015). The age of the Rajang Unconformity is uncertain and is in the range of c. 45 to c. 40 Ma.

altogether, notably those who have advocated South China Sea formation to be a consequence of India indentation of Asia (e.g. Tapponnier *et al.*, 1986; Replumaz & Tapponnier, 2003; Replumaz *et al.*, 2004).

In this paper we explain how the term Proto-South China Sea has been used for oceanic lithosphere subducted at different times under Borneo. We highlight differences in the geological history of different parts of northern Borneo with the aid of a simplified geological map (Figure 3) and stratigraphic chart (Figure 4) for the Late Cretaceous to Quaternary and show how the geology of northern Borneo is linked to different episodes of subduction. We then discuss the evidence for Proto-South China Sea subduction between the Eocene and Early Miocene, and show that the subducted lithosphere can be identified in the mantle below Borneo and the Philippines.

CRETACEOUS TO EOCENE

Stratigraphy

Sarawak has been divided into a number of sub-parallel zones (Figure 5) with different stratigraphic character, which from south to north are the West Borneo Basement (which broadly equates to the Schwaner Mountains of Kalimantan), Kuching Zone, Sibu Zone and Miri Zone (Haile, 1974). The Kuching Zone is composed of fluvial to marginal marine sediments (Figure 4). These include the Kayan Sandstone and Silantek Formation, Tutoop Sandstone (Plateau Sandstone) and Ketungau Formation of the Ketungau Basin in West Sarawak and NW Kalimantan (Haile, 1957; Liechti *et al.*, 1960; Wolfenden & Haile, 1963; Wilford & Kho, 1965; Muller, 1968; Kanno, 1978; Tan, 1979; Heryanto & Jones, 1996) deposited from the Late Cretaceous (Maastrichtian) to the Middle Eocene (Liechti *et al.*, 1960; Tan, 1979; Pieters *et al.*, 1987; Morley, 1998; Breitfeld, 2015).

The boundary between the Kuching Zone and Sibu Zone is the Lupar Line (Haile, 1974). The Sibu Zone (Figure 4) includes sedimentary rocks of similar age to the Kuching Zone, but in contrast is predominantly composed of turbiditic deep marine sediments of the Belaga Formation (Kirk, 1957; Haile, 1957; Liechti et al., 1960; Wolfenden, 1960; Tan, 1979; Hutchison, 1996; Bakar et al., 2007; Galin, 2013), which together with the Lupar Formation (and the Embaluh Formation in Kalimantan) form the Rajang Group (Haile, 1962; Tate, 1991; Hutchison, 1996; Moss, 1998). The metamorphosed turbiditic Mulu Formation of the Miri Zone (Liechti et al., 1960; Haile, 1962) and the metamorphosed turbiditic Sapulut Formation in Sabah (Collenette, 1965) are lateral equivalents and can therefore be grouped within the Rajang Group. In NE Kalimantan the turbiditic deep marine to slope Mentarang and Lurah Formations are potential equivalents of the Rajang Group, while the Long Bawan Formation may include marine successions (BRGM, 1982). However, the three NE Kalimantan formations are not very well studied, and their correlation with Sarawak and Sabah is only poorly understood.

Recent work on heavy mineral and detrital zircon geochronology shows that the Rajang Group is the lateral deep marine equivalent of the fluvial to shallow marine deposits in the Kuching Zone (Galin, 2013; Breitfeld, 2015) and these formed part of a single continental margin sedimentary system.

Rajang Group: accretionary prism?

The deep water and deformed character of much of the Rajang Group led to the suggestion that it represented an accretionary prism (e.g. Tan, 1979, 1982) related to southward subduction at the Lupar Line, which was therefore a former subduction zone. Haile (1973) discussed the Lupar valley and concluded it had some features of a former subduction zone, but many that were missing. He also commented that the Rajang Group contains no chert or ophiolites and is not a melange. Hutchison (1996) interpreted the Rajang Group as a north-facing accretionary prism, related to Late Cretaceous (85 Ma) to Late Eocene (45 Ma) southwards subduction beneath the Kuching Zone, i.e. at the Lupar Line (Figure 6). Hutchison (2010a) included the Belaga Formation of Sarawak, the Embaluh Group, Malinau Formation and Mentarang Formation of Kalimantan and the Sapulut and Trusmadi Formations of Sabah. Hutchison (1996, 2005, 2010a) argued that older parts of the Rajang Group were accretionary, but that subduction ceased in the Paleocene before most of the turbidites were deposited. For the younger turbidites he followed Moss (1998) who suggested they were deposited in a remnant ocean basin. Moss (1998) specifically excluded an accretionary setting and argued that subduction had ceased in the Late Cretaceous based on an absence of southwards dipping thrusts and, southward rather than northward, younging of the Embaluh Group, which unlike much of the other formations in the Rajang Group is not metamorphosed. Williams *et al.* (1988) and Moss (1998) concluded that subduction-related magmatism in the Schwaner Mountains arc had ceased by about 80



Figure 5: The structural zones of NW Borneo simplified after Haile (1974).



Figure 6: Schematic cross sections across Sarawak based on Moss (1998), redrawn and slightly modified from Hutchison (2010a), who interpreted subduction at the Lupar Line and the Rajang Group (Sibu Zone) to have been deposited on Proto-South China Sea crust which had been subducted between 130 and 80 Ma.

Bulletin of the Geological Society of Malaysia, Volume 63, June 2017

Ma, and Hutchison (1996) observed that the subduction history inferred by Tan & Lamy (1990) and Hazebroek & Tan (1993), from Late Cretaceous to Late Eocene, is not marked by subduction-related Eocene volcanic arcs.

Oceanic subduction

According to Hutchison (1996) the Rajang Group was a north-facing accretionary prism (Figure 6), related to Cretaceous to Eocene southwards subduction of the Danau Sea of Haile (1994). He insisted that the Danau Sea was not related to the younger South China Sea and should not be referred to as the Proto-South China Sea. Despite this, he nonetheless showed subduction of a Proto-South China Sea in the Late Cretaceous beneath the Rajang Group (Figure 6). Hutchison (2010a) also shows the Proto-South China Sea subducting southwards under Sarawak in the Early to Late Cretaceous (130-80 Ma), and Hutchison stated that the Sibu Zone was deposited on Proto-South China Sea oceanic crust. Although no maps were provided, Hutchison (2010a) reported that "the northern and southern Sundaland continents continued to converge until collision eliminated the Proto-South China Sea ocean basin, with small remnants uplifted as ophiolite along the Lupar Line. Extinction of the turbidite basin and uplift of the compressed Sibu Zone was interpreted by Hutchison (2005) to have taken place at the end of the Eocene, during the Sarawak Orogeny, discussed further below.

Moss (1998) continued with this usage when he commented on earlier interpretations of ocean crust subducted under Borneo during the Late Cretaceous as "invariably ascribed to subduction of a large proto-South China Sea, with the Schwaner Mountain granites marking the position of the former volcanic arc". In contrast to Hutchison (1996), Moss argued that oceanic subduction beneath Borneo had ceased by about 80 Ma and the Rajang-Embaluh turbidites were deposits of a remnant oceanic basin (Figure 7) and were not an accretionary complex directly



Figure 7: Paleogeography of the Borneo area in the mid Cretaceous as interpreted by Moss (1998) showing unsubducted Proto-South China Sea north of Sabah and trapped oceanic crust of the Proto-South China Sea on which Embaluh–Rajang Group sediments were later deposited.

related to subduction. King *et al.* (2010) appear to be the only other authors who have named crust subducted during the Cretaceous under Borneo as a Proto-South China Sea.

It is clear that both Hutchison (1996), Moss (1998) and later King et al. (2010) used the term Proto-South China Sea in an entirely different way from previous authors. This oceanic crust was subducted beneath Sarawak and NW Kalimantan, but subduction had ceased either during the Cretaceous or by the Late Eocene and was completed long before subduction of oceanic crust began beneath Sabah and Cagayan as proposed by Hinz et al. (1991). We consider it is confusing to call both the Proto-South China Sea. It is also arguable if pre-Cenozoic subduction should be related to the South China Sea in any way; many authors consider Cretaceous subduction under Borneo as part of a Paleo-Pacific subduction margin. We therefore suggest restricting the name Proto-South China Sea and acronym (PSCS) to the slab subducted mainly beneath Sabah and Cagayan between the Eocene and Early Miocene. We call the lithosphere subducted under Borneo in the Cretaceous the Paleo-Pacific Ocean (PPO).

Our view is that the various melanges including ophiolitic rocks (Lubok Antu, Boyan, etc.) are all part of a wide accretionary zone at the eastern margin of Asia and Sundaland which includes fragments of continental crust (e.g. the Semitau block of Metcalfe, 1990; rocks dredged from the Dangerous Grounds reported by Kudrass et al., 1986) that formed during Triassic to Cretaceous subduction of the Paleo-Pacific Ocean, as suggested by Breitfeld et al. (2017) and Hennig et al. (2017). Hall & Sevastjanova (2012) suggested that the Lupar Line is not a collisional suture, but a strike-slip zone separating the emergent area of the Schwaner Mountains, coastal plains and shallow shelf to the south, from a deep-water turbidite basin further north. We consider that displacements on this strike-slip zone exhumed some of the ophiolitic rocks in the Lubok Antu melange, probably between the Late Cretaceous and Eocene.

EOCENE TO MIOCENE

Stratigraphy

The regions in which the Cretaceous and Paleogene subduction episodes occurred are separated by the West Baram Line (Figure 3) which is the position of a major stratigraphic change in the Middle to Late Eocene (Figure 4). To the SW of the West Baram Line deep water sedimentation ceased and the Rajang Group is overlain by terrestrial to shallow marine sediments of the Miri Zone, which are largely undeformed. Hutchison (e.g. 1996, 2005) attributed this change to a collisional event named by him the Sarawak Orogeny.

Sarawak Orogeny

Hutchison (1996) proposed the Sarawak Orogeny deformed the deep water sediments of the Rajang Group in the Late Eocene at c.45 Ma due to collision of a Balingian–Luconia continental block with SW Borneo about

Bulletin of the Geological Society of Malaysia, Volume 63, June 2017

15 Ma earlier, following southwards subduction which terminated by c.60 Ma. The age of the orogeny was based on unconformities described by earlier workers (Wolfenden, 1960; Adams & Haak, 1962; Haile & Ho, 1991). Later, Hutchison (2004, 2005) assigned the orogeny a younger age of c.37 Ma and described it as the result of indentation of a promontory of northern Sundaland following closure of the proto-South China Sea (Hutchison, 2010a). Hall (2012) and Hall & Sevastjanova (2012) have questioned the age and interpretation of the unconformity.

Borneo Geological Survey geologists did recognise an episode of folding in the Late Eocene (Kirk, 1957; Wolfenden, 1960; Haile, 1962), but before 37 Ma, and as one event in a long Cretaceous to Eocene history of deformation. Wolfenden (1960) identified an unconformity older than uppermost Eocene, and Adams & Haak (1962) showed it was older than the East Indies Letter Stage Tb, indicating that the oldest rocks above the unconformity include the oldest part of the Upper Eocene, which would mean the unconformity is older than c. 40 Ma. Haile & Ho (1991) did not date it. In Sarawak, Wolfenden (1960) noted the absence of a marked angular unconformity in some areas and commented that the "stratigraphic evidence is difficult to reconcile with the concept of an upper Eocene orogeny that caused the entire.....Rajang Group to be folded" and observed that "deformation accompanied deposition".

Hall & Sevastjanova (2012) suggested that the Lupar Line is not a collisional suture, but a strike-slip zone, separating the emergent area of the Schwaner Mountains, coastal plains and shallow shelf to the south, from a deep-water turbidite basin further north. Syn-depositional deformation in the deep-water basin occurred between the Late Cretaceous and the Late Eocene. The Sarawak Orogeny was suggested to mark not a collisional event but a regional plate boundary reorganisation that occurred when Australia began to move north at about 45 Ma, initiating subduction on the north side of Borneo (Hall et al., 2008), beneath Sabah and Cagayan, and south of Java (e.g. Hall et al., 2009; Hall, 2002, 2012). The unconformity which terminated the deposition of the Rajang Group and its equivalents in the Middle to Late Eocene is named here the 'Rajang Unconformity' (Figure 3).

Late Eocene change: Sarawak to Sabah

Whatever the cause, there was an important change in Sarawak in the Late Eocene and highly deformed deep water turbidites are unconformably overlain by slightly tilted, but otherwise undeformed, terrestrial/marginal marine sedimentary rocks (Figure 4). In Sarawak, sedimentation resumed in the Late Eocene to Early Oligocene in the Miri Zone and potentially in the Sibu Zone with the fluvial to marginal marine Tatau and Nyalau Formations of Late Eocene/Early Oligocene to Early Miocene age (Kirk, 1957; Wolfenden, 1960; Hassan, 2004; Wong, 2011) SW of the West Baram Line.

NE of the West Baram Line, sedimentation was dominated by deep marine successions. The Setap Shale

is the time equivalent of the Nyalau Formation in the Miri Zone (Haile, 1962; Hutchison, 2005), and the Temburong and Crocker Formations in Brunei and Sabah represent shaly to sand-dominated turbidites (Collenette, 1958, 1965; Wilson & Wong, 1964; van Hattum et al., 2006, 2013). The turbiditic Labang Formation (Collenette, 1965; Haile & Wong, 1965) and the marine Kulapis Formation (Newton-Smith, 1967; Clennell, 1992; Noad, 1998) are the eastern equivalents of the Crocker Formation. An exception to the overall deep marine character is the Upper Eocene to Lower Miocene Melinau Limestone Formation, which is interpreted as a local marine carbonate platform some distance from the shore (Liechti et al., 1960; Adams & Haak, 1962; Adams, 1965). However, Adams & Wilford (1972) described other Oligocene limestone exposures in the Melinau region which they interpreted as deep water limestones.

In Sabah the partially metamorphic Eocene Trusmadi Formation (Collenette, 1958, 1965) could represent the older parts of the Crocker Formation (as shown in Figure 4), but could also be part of the Rajang Group below the unconformity as shown by Hutchison (2005). The Crocker Formation is terminated by the Early Miocene Top Crocker Unconformity (TCU) that separates it from younger fluvial to shallow marine sediments (van Hattum, 2005; Hall *et al.*, 2008; van Hattum *et al.*, 2013).

Sarawak: western limit of Proto-South China Sea subduction

All the early interpretations of subduction under Borneo (Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz *et al.*, 1991) interpreted subduction to have begun beneath Borneo in the Eocene in an area which extended far to the west, into or beyond Sarawak (Figure 2). It is now evident that the westernmost limit of Proto-South China Sea subduction was the West Baram Line. Recent work also now makes clear that grouping together the deep water Rajang Group and deep water Crocker Formation (and associated formations) is unjustified and misleading. During the period when the Crocker Formation was being deposited NE of the West Baram Line there was terrestrial to shelf sedimentation to the SW (Figures 3 and 4).

Although the idea of south-directed subduction was introduced quite early in the 1970s (Hamilton, 1973, 1979; Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz *et al.*, 1991) and incorporated in many later tectonic interpretations of Sabah and Sulu Sea (e.g. Hutchison, 1989; Hazebroek & Tan, 1993; Hall, 1996; Hutchison *et al.*, 2000; Hall & Wilson, 2000) it is worth revisiting the evidence for subduction since both the size and even the existence of the Proto-South China Sea have been challenged in recent years.

Evidence for subduction c. 45-20 Ma

The case for subduction was strongly influenced by Hamilton's early interpretations, particularly his suggestions of subduction melanges. He drew attention to an association of melanges and broken formations, calc-alkaline igneous rocks, cherts, basalts, ophiolite fragments, and serpentinised

peridotites and interpreted them as the result of Early and Middle Cenozoic subduction beneath northern Borneo.

Basement

Northern Borneo has a basement of ophiolitic rocks, but with possible older crust indicated by K-Ar ages of igneous and metamorphic rocks. The Chert-Spilite Formation includes cherts and limestones of Cretaceous age. Metamorphic rocks were described as crystalline basement (Reinhard & Wenk, 1951; Dhonau & Hutchison, 1966; Koopmans, 1967) and most of the protoliths are basic and all could be deformed ophiolitic rocks. Omang & Barber (1996) suggested the crystalline basement rocks are essentially equivalent to the Cretaceous Darvel Bay ophiolite complex (Dhonau & Hutchison, 1966; Hutchison, 1975, 1978). They are intruded by dioritic and granitic rocks which could represent arc plutonic rocks intruded into an older ophiolitic basement. Hall & Wilson (2000) concluded that descriptions by Leong (1974) and earlier workers suggest formation of the crystalline basement in a Mesozoic intra-oceanic arc. There are however, a few rocks described which are of possible continental origin, such as and alusite-garnet-mica and sillimanite-garnet schists (Leong, 1974), although none of these rocks are dated. Fragments of ophiolitic rocks are found as clasts in Eocene sediments, suggesting that the Sabah ophiolite complex had been obducted in either the latest Cretaceous or earliest Paleogene (Omang & Barber, 1996). No Cenozoic ocean floor basalts or deep marine sediments that could represent Paleogene accreted oceanic crust are known.

HP-LT metamorphic rocks

High pressure metamorphic rocks, including HP-LT rocks typical of subduction metamorphism, have been reported from a number of parts of Sabah. None are dated. Eclogites were described from the Dent peninsula area (Reinhard & Wenk, 1951) and Haile & Wong (1965) reported glaucophane-bearing rocks as well as a variety of other metamorphic rocks. Morgan (1974) estimated pressures and temperatures of 850°C and 19 kbar for the garnet pyroxenites from the Dent peninsula indicating upper mantle conditions. Parkinson et al. (1998) suggested these rocks, accompanied by garnet-kyanite amphibolite, eclogite and blueschist, found within Miocene conglomerates, had a subduction origin.

Kirk (1968) mentioned quartz-muscovite-glaucophane schist and glaucophane amphibolites as float in the Labuk River north of Telupid. Johnston & Walls (1974) reported glaucophane-bearing metabasalts from the Telupid area which could be part of the Chert-Spilite Formation or Crocker Formation. Hutchison et al. (2000) mentioned piedmontite and confirmed glaucophane-epidote facies rocks from Telupid. Leong (1978) suggested all these rocks belonged to a Sabah subduction-related blueschist belt.

Crocker and associated formations

Hamilton (1973, 1979) described and illustrated numerous examples of deformed deep-water Eocene and

Oligocene clastic sediments of the Crocker Formation which he interpreted as part of a subduction-imbricated terrain. Subsequent studies have confirmed the deep water turbidite character of the sediments (e.g. Crevello, 2002; van Hattum, 2005, Lambiase et al., 2008; Cullen et al., 2012; van Hattum et al., 2013) and the common synsedimentary deformation. The character of all the deepwater sediments and the syn-depositional deformation of the Crocker Formation and associated formations such as the Temburong, Kulapis and Labang Formations is consistent with an accretionary forearc setting, with elevated forearc highs from which ophiolitic debris was eroded (van Hattum, 2005; van Hattum et al., 2006, 2013) and where limestones (e.g. Gomantong Formation) may have been deposited (McMonagle et al., 2011). However, none of these features requires a subduction-related forearc setting. Rangin et al. (1990a) described the chaotic complexes of southern Sabah as large SE-dipping thrust slices imbricated with a Late Paleogene to Early Neogene volcanic arc sequence, and with large thrust slices of dismembered ophiolites which does suggest an accretionary forearc setting.

The heavy mineral assemblages of the Eocene-Lower Miocene Crocker Formation are dominated by the ultrastable minerals zircon, tourmaline and less abundant rutile, with variable quantities of other minerals which are locally abundant including garnet and apatite (van Hattum, 2005). Chrome spinel is nearly always present, but never exceeds 4% of the total heavy mineral suite. Pyroxene and amphibole are present in a few samples which could indicate a volcanic arc source. No detrital HP-LT minerals have been found so far in the Crocker Formation (van Hattum, 2005). Euhedral unabraded zircons are quite common. However, the youngest zircon in the 4 samples from which detrital zircons have been dated is 73 Ma (van Hattum et al., 2006, 2013) and is therefore much older than the depositional age. The heavy mineral assemblages of the Eocene Trusmadi Formation sandstones resemble those of the Crocker Formation and in the single sample from which detrital zircons have been dated the youngest zircon is 50 Ma indicating Eocene igneous activity. The Oligocene Labang Formation contains some of the least mature sandstones of Sabah, with a possible magmatic arc provenance based on their light and heavy mineral compositions (van Hattum, 2005). Labang Formation sandstones contain abundant volcanic lithic grains. Typical heavy minerals include zircon, tourmaline, apatite, garnet, rutile, chrome spinel, pyroxene and hornblende. The preservation and relative abundance of pyroxene and hornblende suggest a volcanic arc contribution. No detrital zircons from the Labang Formation have yet been dated.

Volcanic rocks

Pre-Neogene volcanic rocks are rather uncommon in west and central Sabah. Basalts are sometimes found intercalated in the Crocker Formation, presumably faulted, although we have not seen contacts, and could represent contemporaneous igneous activity, but could alternatively be tectonically incorporated slices of basement.

An obvious Paleogene volcanic arc is not known. However, the likely position of the arc suggests it could be buried beneath Neogene basins of central and eastern Sabah. Near Sandakan (Figure 1), at such a position, Rangin *et al.* (1990a) reported the mosque is built on a folded sequence of tuffs, greywackes and massive andesitic flows and microbreccias. Late Oligocene nannoplankton (NP25) were found in this sequence, which is covered unconformably by slightly deformed clastic sediments of the Neogene Sandakan Formation. Bergman *et al.* (2000) reported a K-Ar age of 77 ± 4 Ma from plagioclase (they commented this may be anomalous due to very low K content and possible excess ⁴⁰Ar) and an apatite fission track age of 34 ± 8 Ma from an andesite tuff in this sequence.

Detrital zircons in Paleogene sandstones (van Hattum *et al.*, 2006, 2013) indicate a maximum depositional age (MDA) of 56 Ma for the Sapulut Formation and 50 Ma for the Trusmadi Formation, confirming Eocene volcanic activity. No younger ages have been obtained from detrital zircons in the Crocker Formation, although only 4 sandstone samples have been dated. Probable tuff bands are present in the Crocker Formation in Sabah, but none of these volcanic horizons have yet been dated.

Suggate (2011) dated detrital zircons from 11 samples of Neogene sandstones deposited above the TCU of which 7 contain Eocene zircons (41-48 Ma). These could have come from Borneo, but some in the Kudat Formation could come from Palawan (Suggate & Hall, 2013; Suggate *et al.*, 2014). However, one Miocene Tanjong Formation sample from one of the circular basins in Sabah contains a population of Eocene to Early Miocene zircons with ages of 48-22 Ma (Suggate, 2011) which can only have come from a Borneo source indicating Paleogene volcanic activity.

Cagayan Ridge

Hamilton (1979) tentatively suggested the Cagayan Ridge (Figure 1) could be a Paleogene Arc. Hinz et al. (1991) reported that petrological and geochemical data indicates continental crust below Oligocene to Miocene volcanic rocks. Kudrass et al. (1990) dredged samples of andesites and dacites from the northernmost Cagayan Ridge. They reported K-Ar ages of 158-36 Ma from amphiboles which may be inherited xenocrysts, and the oldest plagioclases of 26-22 Ma. However, they also mentioned problems of excess Ar and Ar loss and concluded there was definite Neogene and possibly some older volcanic activity. Rangin & Silver (1991) reviewed all the data from the Cagayan Ridge after ODP Leg 124 drilling. They observed that the oldest rocks dredged at the northern end of the ridge were close to the axis of a major canyon flowing SE from Panay Island, where the pre-Neogene volcanic rocks of the Philippine arc are exposed. Thus, they were confident of Neogene volcanic activity at 14.7 Ma at Sites 769 and 771 (Figure 1), but were cautious about older volcanism, although they suggested the Cagayan Ridge may have extended SW as far as the Oligocene volcaniclastic rocks of the Sandakan area in Sabah.

Discussion

Overall, the case for Paleogene subduction beneath northern Borneo is strong, but not indisputable. The evidence suggests a pre-Cenozoic ophiolitic/arc basement in Sabah, with hints of deep continental crust (Leong, 1974; Macpherson et al., 2010). There is convincing evidence of HP-LT subduction-related metamorphism, but the age is not known and could be Mesozoic. The Crocker and associated formations are typical of a volcanic forearc margin, but many of the features could be explained by a Paleogene deep-water continental margin setting without subduction. Most of the sediment came from the unroofed Cretaceous Schwaner magmatic arc, either directly or indirectly, and the Tin Belt of the Thai-Malay peninsula (van Hattum et al., 2006, 2013). There is good evidence for Paleogene volcanism from outcrops near Sandakan, volcanic lithic grains in the Labang Formation, heavy minerals in some sandstones, and ages of detrital zircons in Neogene sediments. However, given the limited number of samples that have been dated it is clear that more work is required to determine the age and importance of volcanic activity. Given the interpreted size of the Proto-South China Sea (Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz et al., 1991) more evidence of a Paleogene arc might be expected, although the subducted embayment narrowed westwards, which would have reduced the amount of subduction, and an arc could be buried below Neogene sediments. Subduction does not always produce magmatic products (e.g. Macpherson & Hall, 2002) and there are several active subduction zones in the region (North Sulawesi Trench, Philippine Trench) without associated volcanic arcs. Alternatively, modifications to reconstructions could reduce the interpreted size of the Proto-South China Sea (Cullen, 2010).

SEISMIC TOMOGRAPHY

It is now accepted that the mantle contains a record of subduction, because subducted lithosphere produces a strong temperature anomaly which causes slabs to be observed as regions with relatively high seismic velocities. One important test of tectonic models involving subduction is to examine the structure of the mantle, since high-resolution P-wave seismic tomography can reveal such velocity anomalies. Is the Proto-South China Sea identifiable?

Where is the Proto-South China Sea?

Rangin *et al.* (1999b) were the first to consider mantle structure and presented three options based on different tectonic models for the Proto-South China Sea (PSCS): large (Holloway, 1981, 1982; Taylor & Hayes, 1983; Hinz *et al.*, 1991), none (Tapponnier *et al.*, 1986), or small (Rangin *et al.*, 1999b). They pointed out that no large anomaly existed in the position they expected for the PSCS below northern Borneo and the Sulu Sea. This could mean the PSCS had not existed or a high velocity anomaly was not obvious because the PSCS had been small. Alternatively, they proposed that the PSCS was north of its expected position because of clockwise rotation of Borneo, speculated to

have been initiated 10 to 15 million years ago and inferred from GPS measurements (Rangin *et al.*, 1999a) made between 1994 and 1996. They concluded that the PSCS could be identified as a high velocity anomaly present beneath the present South China Sea at depths between 400 and 660 km.

Tang & Zheng (2013) interpreted regional surface wave tomography to show a Proto-South China Sea slab of 500 km length in the upper mantle beneath northern Borneo which is longer than estimated by Rangin *et al.* (1999b), but smaller than interpreted by the Hall (1996, 2002, 2012) reconstructions.

Cullen (2010) also questioned the evidence for a PSCS slab. He reported that P-wave tomographic images of Hafkenscheid (2004) show a high velocity anomaly between 200 km and 300 km under northern Borneo, but concluded it was difficult to relate this to subduction under NW Borneo. He also observed that tomographic images in Replumaz et al. (2004) do not show this anomaly and concluded that although these do not preclude the presence of a subducted slab beneath NW Borneo (since other supposed subduction zones are not imaged) they indicate a relatively small Proto-South China Sea. However, because of its relatively aseismic character the northern Borneo region has poor resolution using P-wave tomography. This could account for the differences between the Hafkenscheid (2004) and Replumaz et al. (2004) images, both of which are based on Utrecht P-wave tomographic models of Spakman. Resolution tests (Hall & Spakman, 2015) show the high velocity anomaly between 200 km and 300 km could be an artefact.

But the lack of resolution in P-wave tomographic models for Borneo is not the main problem. There is a more compelling reason why those searching for the PSCS in the mantle have not found it; they have been looking in the wrong place.

Hall & Spakman (2015) reviewed mantle structure beneath SE Asia using a significantly improved P-wave tomographic model compared to those of previous authors. The tomographic images record subduction beneath the SE Asian region to depths of approximately 1600 km. In the lower mantle a broad and deep anomaly imaged in earlier models below SE Asia now can be seen to have a resolvable internal structure. Upper mantle anomalies beneath SE Asia typically record subduction during the last 10 to 25 Ma (Hall & Spakman, 2015) and the PSCS was subducted before 20 Ma. Thus, if it is not an artefact, the shallow anomaly discussed by Cullen (2010) below North Borneo is young and cannot be due to subduction of the PSCS. Secondly, Borneo and the SE margin of Sundaland have rotated since the Early Miocene (Hall, 1996, 2002, 2012) and therefore the upper mantle directly beneath Borneo is not the place to look. In contrast, in the lower mantle there is an anomaly north of the equator (Hall & Spakman, 2015) that can be traced from below central Borneo with a SW-NE strike to 15°N under the central Philippines (Figure 8A). The tomographic images show an anomaly now at depths below 700 km, with a lateral extent consistent with a wide

Paleogene Proto-South China Sea, which narrowed and terminated towards the SW, and a position consistent with a large Neogene rotation of Borneo. Hall & Spakman (2015) suggest this anomaly is the Proto-South China Sea which subducted southwards from 45 to 20 Ma beneath Sabah and Cagavan consistent with tectonic models (Figure 8B). Figure 8B shows essentially the tectonic model originally proposed by Holloway (1981, 1982), Taylor & Hayes (1983) and Hinz et al. (1991), but with a later rotation of Borneo, based on subsequent palaeomagnetic work (reviewed by Fuller et al., 1999) as discussed by Hall et al. (2008). The other principal difference is that the early models showed subduction of the Proto-South China Sea beneath Sarawak from the Eocene to the Middle Miocene whereas we suggest there was no subduction SW of the West Baram Line beneath Sarawak during this interval (Figure 8B).

NEOGENE

In the Early Miocene the attenuated South China continental margin collided with Sabah and Cagayan terminating subduction, an event which Hutchison (1996) named the Sabah Orogeny. Collision completely eliminated the Proto-South China Sea. SW of the West Baram Line in Sarawak there was little or no deformation. NE of the West Baram Line, the Sabah Orogeny deformed and elevated much of Sabah, and probably Palawan, and produced a major regional unconformity (Figure 4), the Top Crocker Unconformity (TCU of van Hattum *et al.*, 2006, 2013).

The Crocker Formation in western Sabah is unconformably overlain by the Lower Miocene shallow marine Meligan Formation (Wilson & Wong, 1964) and Kudat Formation. To the east of the Crocker Mountains during the Middle and Late Miocene there was subsidence and deposition of a thick succession of fluvio-marine deposits (Noad, 1998; Balaguru *et al.*, 2003; Balaguru & Nichols, 2004) in the Central Sabah Basin (Hutchison, 1992) whose remnants are now found in the circular basins of eastern Sabah (Tongkul, 1993; Clennell, 1996; Balaguru *et al.*, 2003; Tongkul & Chang, 2003).

The collision caused emergence of much of Sabah and the present central highlands of northern Borneo, with folding and thrusting of both basement and cover. However, by the end of the Early Miocene much of present-day Sabah was below or close to sea level (Noad, 1998; Balaguru *et al.*, 2003; Hall *et al.*, 2008) with probably a low elevated range of hills at the position of the Crocker Mountains. Because there are few Neogene rocks in western Sabah this is uncertain, but in Brunei and Sarawak Neogene rocks are preserved and the shelf edge moved broadly northwestwards from the Middle Miocene onwards (e.g. Sandal, 1996; Hazebroek & Tan, 1993; Hutchison, 2005; Cullen, 2010), suggesting a gradual rise and widening of the Crocker Mountains during the Middle and Late Miocene.

To the west of the shelf edge in offshore Brunei and Sabah, and NE of the West Baram Line, is a thick Middle Miocene to Recent deep-water clastic wedge that thins northwards. This is deformed by folds and thrusts to form



Figure 8: A. Lower mantle tomographic depth slice at 800 km from Hall & Spakman (2015) showing high velocity anomalies. Anomalies east of about 110°E are interpreted as the result of subduction since 45 Ma. The anomaly due to the subducted Proto-South China Sea is labelled. B. Reconstruction at 30 Ma modified from Hall (2012) showing interpreted subduction zones. Sunda-Java trench subduction, Proto-South China Sea subduction, subduction below North Sulawesi-East Philippines, and subduction below NE side of Philippine Sea plate all fit well with position of high velocity anomalies in lower mantle. The area shaded brown in Sabah represents the Crocker and related formations on land.

a wide and active fold and thrust belt between the coast and the NW Borneo Trough (Tan & Lamy, 1990; Hazebroek & Tan, 1993; Hinz *et al.*, 1989; McGilvery & Cook, 2003; Morley *et al.*, 2003; Ingram *et al.*, 2004; Franke *et al.*, 2008, 2011; Morley & Leong, 2008; Hesse *et al.*, 2009; Gartrell *et al.*, 2011).

PRESENT

The NW Borneo–Palawan Trough and offshore Sabah fold and thrust belt are often interpreted as features *Bulletin of the Geological Society of Malaysia, Volume 63, June 2017*

resulting from collision, regional compression or subduction. However, there is no seismicity, dipping slab or volcanicity indicating subduction, nor obvious causes of compression.

The area around northern Borneo is largely free of seismicity (Hamilton, 1979; Engdahl *et al.*, 1998). All hypocentres close to Borneo are shallow, with depths less than 50 km, and a majority are shallower than 30 km. Earthquakes for which there are moment tensor solutions show an inconsistent pattern (Simons *et al.*, 2007). In the SE, beneath the Celebes Sea and Mangkalihat peninsula,

most solutions indicate NE-SW compression, whereas those further north, mainly beneath the northeast Sabah coast and the Sulu Sea, suggest predominantly NW-SE extension and some strike-slip movements.

GPS observations (Simons *et al.*, 2007) indicate active deformation of northern Borneo with small displacements relative to Sundaland and southern Borneo, suggesting broadly SW-directed movements up to 8 mm/yr. It is doubtful if they record anything other than movements of the upper crust. The orientation of GPS vectors is different from motions indicated by earthquake solutions and suggested by the orientation of structures offshore (e.g. Hazebroek & Tan, 1993; Hutchison, 2005; Hesse *et al.*, 2009; Morley, 2009), and most GPS stations are close to the coast in areas likely to be underlain by weak sedimentary rocks or unconsolidated sediments; GPS vectors could simply reflect shallow displacements related to gravity-driven movements.

There are no active volcanoes in Sabah nor evidence for magmatism related to possible subduction at the troughs. In the Dent and Semporna peninsulas Mio-Pliocene volcanism was related to northward subduction of the Celebes Sea. Plio-Pleistocene volcanism in the Tawau area and Semporna peninsula (Kirk, 1968), and on small islands in Darvel Bay have an ocean-island basalt character that reflects upwelling of melts in the upper mantle into lithospheric thin spots produced during earlier subduction (Macpherson *et al.*, 2010).

The NW Borneo trough is a flexural response to gravitydriven deformation of the sediment wedge, caused by uplift on land that resulted from extension, with a contribution of deep crustal flow as explained by Hall (2013). For most of the offshore foldbelt extension and contraction can be balanced, consistent with a gravity-driven mechanism (Hesse *et al.*, 2009). Further north, the Palawan Trough is the continent-ocean transition (Franke *et al.*, 2011) at the SE edge of the modern South China Sea which formed in the Oligocene by rifting and is clearly unrelated to subduction.

CONCLUSIONS

The term Proto-South China Sea was introduced to describe oceanic crust that formerly occupied the area south of Indochina and South China where the modern South China Sea is situated. The oceanic crust had a Mesozoic age, and has been completely eliminated by subduction beneath northern Borneo and Cagayan. Subduction began in the Eocene and ceased in the Early Miocene. The term Proto-South China Sea and the acronym (PSCS) should be used only for the slab subducted mainly beneath Sabah and Cagayan between the Eocene and Early Miocene. Oceanic crust subducted during earlier episodes of subduction in other areas should be named differently and we name lithosphere subducted under Borneo in the Cretaceous the Paleo-Pacific Ocean (PPO).

There is good evidence for subduction between the Eocene and Early Miocene in Sabah, and the western limit of Proto-South China Sea subduction was the West Baram Line. The subducted Proto-South China Sea can be imaged in the lower mantle using P-wave tomography. There was no subduction beneath Sarawak, SW of the West Baram Line, between the Eocene and Early Miocene where there was terrestrial to marginal marine deposition at this time.

The present-day NW Borneo Trough and Palawan Trough are not subduction trenches. The NW Borneo Trough is a flexural depression due to gravity-driven deformation of the Neogene sediment wedge west of Sabah, with a contribution of deep crustal flow. The Palawan Trough is the continent-ocean transition at the SE edge of the modern South China Sea.

ACKNOWLEDGEMENTS

This work has been supported mainly by the SE Asia Research Group at Royal Holloway, funded by a consortium of oil companies. Financial support has also been provided at times for our work in Borneo by the Royal Society and the London University Central Research Fund. We thank Ian Metcalfe and Tony Barber for helpful comments on the manuscript.

REFERENCES

- Adams, C.G., 1965. The foraminifera and stratigraphy of the Melinau Limestone, Sarawak, and its importance in Tertiary correlation. Quarterly Journal of the Geological Society of London, 121, 283-338.
- Adams, C.G. & Haak, R., 1962. The stratigraphical succession in the Batu Gading area, Middle Baram, North Sarawak. In: Haile, N.S. (Ed.), The Geology and Mineral Resources of the Suai-Baram area, North Sarawak. Geological Survey Department, British Territories in Borneo, Memoir 13, 141-150.
- Adams, C.G. & Wilford, G.E., 1972. On the age and origin of the Keramit and Selidong limestones, Sarawak, East Malaysia. Geological Survey of Malaysia, Geological Papers 1, 28-42.
- Bakar, Z.A.A., Madon, M. & Muhamad, A.J., 2007. Deep-marine sedimentary facies in the Belaga Formation (Cretaceous-Eocene), Sarawak: Observations from new outcrops in the Sibu and Tatau areas. Bulletin of the Geological Society of Malaysia, 53, 35–45.
- Balaguru, A. & Nichols, G.J., 2004. Tertiary stratigraphy and basin evolution, southern Sabah (Malaysian Borneo). Journal of Asian Earth Sciences, 23, 537-5554.
- Balaguru, A., Nichols, G.J. & Hall, R., 2003. The origin of the circular basins of Sabah, Malaysia. Bulletin of the Geological Society of Malaysia, 46, 335-351.
- Barckhausen, U. & Roeser, H.A., 2004. Seafloor spreading anomalies in the South China Sea revisited. In: Clift, P., Wang, P., Kuhnt, W. & Hayes, D.E. (Eds.), Continent-Ocean Interactions within the East Asian Marginal Seas. AGU Geophysical Monograph, 149, American Geophysical Union, 121-125.
- Barckhausen, U., Engels, M., Franke, D., Ladage, S. & Pubellier, M., 2014. Evolution of the South China Sea: Revised ages for breakup and seafloor spreading. Marine and Petroleum Geology, 58, 599-611.
- Barckhausen, U., Engels, M., Franke, D., Ladage, S. & Pubellier, M., 2015. Reply to Chang *et al.*, 2014, Evolution of the South China Sea: Revised ages for breakup and seafloor spreading. Marine and Petroleum Geology, 59, 679-681.
- Ben-Avraham, Z. & Uyeda, S., 1973. The evolution of the China Basin and the Mesozoic palaeogeography of Borneo. Earth and Planetary Science Letters, 18, 365-376.

Bulletin of the Geological Society of Malaysia, Volume 63, June 2017

- Bergman, S.C., Hutchison, C.S., Swauger, D.A. & Graves, J.E., 2000. K:Ar ages and geochemistry of the Sabah Cenozoic volcanic rocks. Bulletin of the Geological Society of Malaysia, 44, 165-171.
- Breitfeld, H.T., 2015. Provenance, stratigraphy and tectonic history of Mesozoic to Cenozoic sedimentary rocks of West and Central Sarawak, Malaysia. PhD Thesis, Royal Holloway University of London. 808 p.
- Breitfeld, H.T., Hall, R., Galin, T., Forster, M.A. & BouDagher-Fadel, M.K., 2017. A Triassic to Cretaceous Sundaland–Pacific subduction margin in West Sarawak, Borneo. Tectonophysics, 694, 35-56.
- BRGM, 1982. Geological mapping and mineral exploration in northeast Kalimantan, 1979–1982. Final report, Final report Rapport du Bureau de Recherches Géologique et Minières 82 RDM 007AO, 120 p.
- Briais, A., Patriat, P. & Tapponnier, P., 1993. Updated interpretation of magnetic anomalies and sea floor spreading stages in the South China Sea: Implications for the Tertiary tectonics of Southeast Asia. Journal of Geophysical Research, 98, 6299-6328.
- Chang, J.-H., Lee, T.-Y., Hsu, H.-H. & Liu, C.-S., 2015. Comment on Barckhausen *et al.*, 2014 – Evolution of the South China Sea: Revised ages for breakup and seafloor spreading. Marine and Petroleum Geology, 59, 676-678.
- Clennell, B., 1992. The melanges of Sabah, Malaysia. PhD Thesis, University of London, London. 483 p.
- Clennell, B., 1996. Far-field and gravity tectonics in Miocene basins of Sabah, Malaysia. In: Hall, R. & Blundell, D.J. (Eds.), Tectonic Evolution of SE Asia. Geological Society of London Special Publication, 106, 307-320.
- Clift, P., Lee, G.H., Duc, N.A., Barckhausen, U., Long, H.V. & Zhen, S., 2008. Seismic reflection evidence for a Dangerous Grounds miniplate: No extrusion origin for the South China Sea. Tectonics, 27, doi:10.1029/2007TC002216.
- Collenette, P., 1958. The geology and mineral resources of the Jesselton-Kinabalu area, North Borneo. Malaysia Geological Survey Borneo Region, Memoir 6. 194 p.
- Collenette, P., 1965. The geology and mineral resources of the Pensiangan and Upper Kinabatangan area, Sabah. Malaysia Geological Survey Borneo Region, Memoir 12. 150 p.
- Crevello, P.D., 2002. The Great Crocker submarine fan: a world class foredeep turbidite system. Proceedings Indonesian Petroleum Association 28th Annual Convention, 377-407.
- Cullen, A., 2010. Transverse segmentation of the Baram-Balabac Basin, NW Borneo: refining the model of Borneo's tectonic evolution. Petroleum Geoscience, 16, 3-29.
- Cullen, A., 2014. Nature and significance of the West Baram and Tinjar Lines, NW Borneo. Marine and Petroleum Geology, 51, 197-209.
- Cullen, A.B., Zechmeister, M.S., Elmore, R.D. & Pannalal, S.J., 2012. Paleomagnetism of the Crocker Formation, northwest Borneo: Implications for late Cenozoic tectonics. Geosphere, 8, 1146-1169.
- Dhonau, T.J. & Hutchison, C.S., 1966. The Darvel Bay area, East Sabah, Malaysia. Malaysia Geological Survey Borneo Region, Annual Report for 1965, 141-160.
- Doutch, H.F., 1992. Aspects of the structural histories of the Tertiary sedimentary basins of East, Central and West Kalimantan and their margins. BMR Journal of Australian Geology and Geophysics, 13, 237-250.
- Engdahl, E.R., van der Hilst, R. & Buland, R., 1998. Global teleseismic earthquake relocation with improved travel times and procedures for depth determination. Bulletin of the

Bulletin of the Geological Society of Malaysia, Volume 63, June 2017

Seismological Society of America, 88, 722-743.

- Franke, D., Barckhausen, U., Heyde, I., Tingay, M. & Ramli, N., 2008. Seismic images of a collision zone offshore NW Sabah/ Borneo. Marine and Petroleum Geology, 25, 606-624.
- Franke, D., Barckhausen, U., Baristeas, N., Engels, M., Ladage, S., Lutz, R., Montano, J., Pellejera, N., Ramos, E.G. & Schnabel, M., 2011. The continent-ocean transition at the southeastern margin of the South China Sea. Marine and Petroleum Geology, 28, 1187-1204.
- Fuller, M., Ali, J.R., Moss, S.J., Frost, G.M., Richter, B. & Mahfi, A., 1999. Paleomagnetism of Borneo. Journal of Asian Earth Sciences, 17, 3-24.
- Galin, T., 2013. Provenance of the deep marine Belaga Formation in the Sibu Zone north of the Lupar Line, Sarawak, Malaysia. MSc Thesis, Royal Holloway University of London. 161 p.
- Gartrell, A., Torres, J. & Hoggmascall, N., 2011. Aregional approach to understanding basin evolution and play systematic in Brunei - unearthing new opportunities in a mature basin. In: International Petroleum Technology Conference, pp. IPTC 1571, 1-5, Bangkok, Thailand.
- Gatinsky, Y.G. & Hutchison, C.S., 1986. Cathaysia, Gondwanaland and the Paleotethys in the evolution of continental Southeast Asia. Bulletin of the Geological Society of Malaysia, 20, 179-199.
- Gebco, 2003. IHO-UNESCO, General Bathymetric Chart of the Oceans, Digital Edition 2003, http://www.gebco.net/.
- Hafkenscheid, E., 2004. Subduction history of the Tethys Oceans reconstructed from plate kinematics and mantle tomography. PhD Thesis, Utrecht University, 208 p.
- Haile, N.S., 1957. The geology and mineral resources of the Lupar and Saribas Valleys, West Sarawak. Malaysia Geological Survey Borneo Region, Memoir 5, 123 p.
- Haile, N.S., 1962. The geology and mineral resources of the Suai-Baram area, north Sarawak, Geological Survey Department British Territories in Borneo, Memoir 13, 176 p.
- Haile, N.S., 1968. The northwest Borneo geosyncline in its geotectonic setting. Bulletin of the Geological Society of Malaysia, Studies in Malaysian Geology 1, 59-60.
- Haile, N.S., 1973. The recognition of former subduction zones in Southeast Asia. In: Tarling, D.H. & Runcorn, S.K. (Eds.), Implications of continental drift to the Earth Sciences, 2, Academic Press, London, 885-891.
- Haile, N.S., 1974. Borneo. In: Spencer, A.M. (Ed.), Mesozoic-Cenozoic Orogenic Belts. Geological Society of London Special Publication, 4, 333-347.
- Haile, N.S., 1994. The Rajang accretionary prism and the Trans-Borneo Danau Suture, Abstract: Tectonic Evolution of SEAsia, December 1994. Geological Society London, 17.
- Haile, N.S. & Ho, C.K., 1991. Geological Field Guide: Sibu-Miri traverse, Sarawak (24 September - 1 October 1991). Petronas Petroleum Research Institute, 37 p.
- Haile, N.S. & Wong, N.P.Y., 1965. The geology and mineral resources of the Dent Peninsula, Sabah. Malaysia Geological Survey Borneo Region Memoir, 16, 199 p.
- Hall, R., 1996. Reconstructing Cenozoic SE Asia. In: Hall, R. & Blundell, D. J. (Eds.), Tectonic Evolution of SEAsia. Geological Society London Special Publication, 106, 153-184.
- Hall, R., 2002. Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. Journal of Asian Earth Sciences, 20, 353-434.
- Hall, R., 2012. Late Jurassic–Cenozoic reconstructions of the Indonesian region and the Indian Ocean. Tectonophysics,

570, 1-41.

- Hall, R., 2013. Contraction and extension in northern Borneo driven by subduction rollback. Journal of Asian Earth Sciences, 76, 399-411.
- Hall, R. & Sevastjanova, I., 2012. Australian crust in Indonesia. Australian Journal of Earth Sciences, 59, 827-844.
- Hall, R. & Spakman, W., 2015. Mantle structure and tectonic history of SE Asia. Tectonophysics, 658, 14-45.
- Hall, R. & Wilson, M.E.J., 2000. Neogene sutures in eastern Indonesia. Journal of Asian Earth Sciences, 18, 787-814.
- Hall, R., van Hattum, M.W.A. & Spakman, W., 2008. Impact of India-Asia collision on SE Asia: the record in Borneo. Tectonophysics, 451, 366-389.
- Hall, R., Clements, B. & Smyth, H.R., 2009. Sundaland: Basement character, structure and plate tectonic development. Proceedings Indonesian Petroleum Association 33rd Annual Convention, IPA09-G-134, 1-27.
- Hamilton, W., 1973. Tectonics of the Indonesian region. Bulletin of the Geological Society of Malaysia, 6, 3-11.
- Hamilton, W., 1979. Tectonics of the Indonesian region. USGS Professional Paper, 1078, 345 p.
- Hassan, K., 2004. Cenozoic. In: Peng, L.C., Leman, S.L., Hassan, K., Nasib, B.M. & Karim, R. (Eds.), Stratigraphic Lexicon of Malaysia. Geological Society of Malyasia, 124-125.
- Hazebroek, H. P. & Tan, D.N.K., 1993. Tertiary tectonic evolution of the NW Sabah Continental Margin. Bulletin of the Geological Society of Malaysia, 33, 195-210.
- Heng, Y.E., 1992. Geological Map of Sarawak, 1:500,000. Geological Survey of Malaysia.
- Hennig, J., Breitfeld, H.T., Hall, R. & Nugraha, A.M.S., 2017. The Mesozoic tectono-magmatic evolution at the Paleo-Pacific subduction zone in West Borneo. Gondwana Research, 48, 292-310.
- Heryanto, R. & Jones, B.G., 1996. Tectonic development of Melawi and Ketungau Basins, Western Kalimantan, Indonesia. Bulletin Geological Research and Development Centre Bandung, 19, 151-179.
- Hesse, S., Back, S. & Franke, D., 2009. The deep-water foldand-thrust belt offshore NW Borneo: Gravity-driven versus basement-driven shortening. Geological Society of America Bulletin, 121, 939-953.
- Hinz, K., Frisch, J., Kempter, E.H.K., Manaf Mohammad, A., Meyer, J., Mohamed, D., Vosberg, H., Weber, J. & Benavidez, J., 1989. Thrust tectonics along the continental margin of Sabah, Northwest Borneo. Geologische Rundschau, 78, 705-730.
- Hinz, K., Block, M., Kudrass, H.R. & Meyer, H., 1991. Structural elements of the Sulu Sea, Phillipines. Geologisches Jahrbuch, Reihe A 127, 483-506.
- Holloway, N.H., 1981. The North Palawan block, Philippines: its relation to the Asian mainland and its role in the evolution of the South China Sea. Bulletin of the Geological Society of Malaysia, 14, 19-58.
- Holloway, N.H., 1982. North Palawan block, Philippines its relation to Asian mainland and role in evolution of South China Sea. American Association of Petroleum Geologists Bulletin, 66, 1355-1383.
- Hutchison, C.S., 1975. Ophiolites in southeast Asia. Geological Society of America Bulletin, 86, 797-806.
- Hutchison, C.S., 1978. Ophiolite metamorphism in northeast Borneo. Lithos, 11, 195-208.
- Hutchison, C.S., 1989. Geological Evolution of South-East Asia. Clarendon Press, Oxford. 376 p.
- Hutchison, C.S., 1992. The Southeast Sulu Sea, a Neogene marginal

basin with outcropping extensions in Sabah. Bulletin of the Geological Society of Malaysia, 32, 89-108.

- Hutchison, C.S., 1996. The 'Rajang Accretionary Prism' and 'Lupar Line' problem of Borneo. In: Hall, R. & Blundell, D.J. (Eds.), Tectonic Evolution of SE Asia. Geological Society London Special Publication, 106, 247-261.
- Hutchison, C.S., 2004. Marginal basin evolution: the southern South China Sea. Marine and Petroleum Geology, 21, 1129–1148.
- Hutchison, C.S., 2005. Geology of North-West Borneo. Elsevier, Amsterdam. 421 p.
- Hutchison, C.S., 2010a. Oroclines and paleomagnetism in Borneo and South-East Asia. Tectonophysics, 496, 53-67.
- Hutchison, C.S., 2010b. The North-West Borneo Trough. Marine Geology, 271, 32-43.
- Hutchison, C.S., Bergman, S.C., Swauger, D.A. & Graves, J.E., 2000. A Miocene collisional belt in north Borneo: uplift mechanism and isostatic adjustment quantified by thermochronology. Journal of the Geological Society London, 157, 783-793.
- Ingram, G.M., Chisholm, T.J., Grant, C.J., Hedlund, C.A., Stuart-Smith, P. & Teasdale, J., 2004. Deepwater North West Borneo: hydrocarbon accumulation in an active fold and thrust belt. Marine and Petroleum Geology, 21, 879-887.
- Johnston, J.C. & Walls, P.J., 1975. Geology of the Telupid area, Sabah. Geological Survey of Malaysia, Annual Report for 1973, 213-220.
- Kanno, S., 1978. Brackish molluscan fauna (Upper Eocene) from the Silantek Formation in West Sarawak, Malaysia. Geology and Palaeontology of S.E. Asia (University of Tokyo Press), 20, 103-112.
- Karig, D.E., 1971. Origin and development of marginal basins in the Western Pacific. Journal of Geophysical Research, 76, 2542-2561.
- King, R.C., Tingay, M.R.P., Hillis, R.R., Morley, C.K. & Clark, J., 2010. Present-day stress orientations and tectonic provinces of the NW Borneo collisional margin. Journal of Geophysical Research, 115, B10415, doi:10.1029/2009JB006997.
- Kirk, H.J.C., 1957. The Geology and Mineral Resources of the Upper Rajang and adjacent areas. British Territories Borneo Region Geological Survey, Memoir 8, 181 p.
- Kirk, H.J.C., 1968. The igneous rocks of Sarawak and Sabah. Geological Survey of Malaysia, Borneo Region Bulletin, 5, 210 p.
- Koopmans, B.N., 1967. Deformation of the metamorphic rocks and the chert spilite formation in the southern part of the Darvel Bay area, Sabah. Geological Survey of Malaysia, Borneo Region Bulletin, 8, 14-24.
- Kudrass, H.R., Wiedecke, M., Cepek, P., Kreuzer, H. & Muller, P., 1986. Mesozoic and Cainozoic rocks dredged from the South China Sea (Reed Bank area) and Sulu Sea and their significance for plate-tectonic reconstructions. Marine and Petroleum Geology, 3, 19-30.
- Kudrass, H.R., Müller, P., Kreuzer, H. & Weiss, W., 1990. Volcanic rocks and Tertiary carbonates dredged from the Cagayan Ridge and the Southwest Sulu Sea, Philippines. In: Silver, E.A., Rangin, C., von Breymann, M.T. *et al.* (Eds.), Proceedings of the Ocean Drilling Program, Initial Reports 124, 93-100.
- Lambiase, J.J., Tzong, T.Y., William, A.G., Bidgood, M.D., Brenac, P. & Cullen, A.B., 2008. The West Crocker formation of northwest Borneo: A Paleogene accretionary prism. In: Draut, A.E., Clift, P.D. & Scholl, D.W. (Eds.), Formation and Applications of the Sedimentary Record in Arc Collision Zones. Geological Society of America Special Paper, 436, 171-184.
- Lee, T.-Y. & Lawver, L.A., 1994. Cenozoic plate reconstructions

of the South China Sea region. Tectonophysics, 235, 149-180.

- Lee, T.-Y. & Lawver, L.A., 1995. Cenozoic plate reconstruction of Southeast Asia. Tectonophysics, 251, 85-138.
- Leong, K.M., 1974. The geology and mineral resources of the Upper Segama Valley and Darvel area, Sabah, Malaysia. Geological Society of Malaysia Memoir, 4, 354 p.
- Leong, K.M., 1978. The 'Sabah Blueschist Belt' A preliminary note. Warta Geologi, Geological Society of Malaysia Newsletter, 4, 45-51.
- Li, C.-F., Xu, X., Lin, J., Sun, Z., Zhu, J., Yao, Y., Zhao, X., Liu, Q., Kulhanek, D.K., Wang, J., Song, T., Zhao, J., Qiu, N., Guan, Y., Zhou, Z., Williams, T., Bao, R., Briais, A., Brown, E.A., Chen, Y., Clift, P.D., Colwell, F.S., Dadd, K.A., Ding, W., Almeida, I.H., Huang, X.-L., Hyun, S., Jiang, T., Koppers, A.A.P., Li, Q., Liu, C., Liu, Z., Nagai, R.H., Peleo-Alampay, A., Su, X., Tejada, M.L.G., Trinh, H.S., Yeh, Y.-C., Zhang, C., Zhang, F. & Zhang, G.-L., 2014. Ages and magnetic structures of the South China Sea constrained by deep tow magnetic surveys and IODP Expedition 349. Geochemistry, Geophysics, Geosystems, 15, 4958-4983.
- Li, C.-F., Li, J., Ding, W., Franke, D., Yao, Y., Shi, H., Pang, X., Cao, Y., Lin, J., Kulhanek, D.K., Williams, T., Bao, R., Briais, A., Brown, E.A., Chen, Y., Clift, P.D., Colwell, F.S., Dadd, K.A., Hernández-Almeida, I., Huang, X.-L., Hyun, S., Jiang, T., Koppers, A.A.P., Li, Q., Liu, C., Liu, Q., Liu, Z., Nagai, R.H., Peleo-Alampay, A., Su, X., Sun, Z., Tejada, M.L.G., Trinh, H.S., Yeh, Y.-C., Zhang, C., Zhang, F., Zhang, G.-L. & Zhao, X.C.J.B., 2015. Seismic stratigraphy of the central South China Sea basin and implications for neotectonics. Journal of Geophysical Research: Solid Earth, 120, 1377-1399.
- Liechti, P., Roe, F.W. & Haile, N.S., 1960. The Geology of Sarawak, Brunei and the western part of North Borneo. Geological Survey Department, British Territories of Borneo, Bulletin 3, 360 p.
- Lim, P.S. & Heng, Y.E., 1985. Geological Map of Sabah 1:500,000, Geology Survey of Malaysia.
- Macpherson, C.G. & Hall, R., 2002. Timing and tectonic controls on magmatism and ore generation in an evolving orogen: evidence from Southeast Asia and the western Pacific. In: Blundell, D.J., Neubauer, F. & von Quadt, A. (Eds.), The timing and location of major ore deposits in an evolving orogen. Geological Society of London Special Publication, 204, 49-67.
- Macpherson, C.G., Chiang, K.K., Hall, R., Nowell, G.M., Castillo, P.R. & Thirlwall, M.F., 2010. Plio-Pleistocene intra-plate magmatism from the southern Sulu Arc, Semporna peninsula, Sabah, Borneo: Implications for high-Nb basalt in subduction zones. Journal of Volcanology and Geothermal Research, 190, 25-38.
- McGilvery, T.A. & Cook, D.L., 2003. The influence of local gradients on accommodation space and linked depositional elements across a stepped slope profile, offshore Brunei. In: Roberts, H.R., Rosen, N.C., Fillon, R.F. & Anderson, J.B. (Eds.), Shelf Margin Deltas and Linked Down Slope Petroleum Systems: Global Significance and Future Exploration Potential. GCSSEPM (Gulf Coast Section of the SEPM), 23rd Annual GCSSEPM Foundation Bob F. Perkins Research Conference, 387-419.
- McMonagle, L.B., Lunt, P., Wilson, M.E.J., Johnson, K.G., Manning, C. & Young, J., 2011. A re-assessment of age dating of fossiliferous limestones in eastern Sabah, Borneo: Implications for understanding the origins of the Indo-Pacific marine biodiversity hotspot. Palaeogeography, Palaeoclimatology, Palaeoecology, 305, 28-42.
- Metcalfe, I., 1990. Allochthonous terrane processes in Southeast Asia. Philosophical Transactions Royal Society of London,

A331, 625-640.

- Morgan, B.A., 1974. Chemistry and mineralogy of garnet pyroxenites from Sabah, Malaysia. Contributions to Mineralogy and Petrology, 48, 301-314.
- Morley, C.K. 2009. Growth of folds in a deep-water setting. Geosphere, 5, 59-89.
- Morley, C.K. & Leong, L.C., 2008. Evolution of deep-water synkinematic sedimentation in a piggyback basin, determined from three-dimensional seismic reflection data. Geosphere, 4, 939-962.
- Morley, C.K., Back, S., Crevello, P., van Rensbergen, P. & Lambiase, J.J., 2003. Characteristics of repeated, detached, Miocene– Pliocene tectonic inversion events, in a large delta province on an active margin, Brunei Darussalam, Borneo. Journal of Structural Geology, 25, 1147-1169.
- Morley, R.J., 1998. Palynological evidence for Tertiary plant dispersals in the SE Asian region in relation to plate tectonics and climate. In: Hall, R. & Holloway, J. D. (Eds.), Biogeography and Geological Evolution of SE Asia. Backhuys Publishers, Leiden, The Netherlands, 211-234.
- Moss, S.J., 1998. Embaluh Group turbidites in Kalimantan: evolution of a remnant oceanic basin in Borneo during the Late Cretaceous to Palaeogene. Journal of the Geological Society London, 155, 509-524.
- Muller, J., 1968. Palynology of the Pedawan and Plateau Sandstone Formations (Cretaceous-Eocene) in Sarawak, Malaysia. Micropaleontology, 14, 1-37.
- Newton-Smith, J., 1967. Bidu-Bidu Hills area, Sabah, East Malaysia, Borneo Region Malaysia Geological Survey Report 4, 109 p.
- Noad, J.J., 1998. The sedimentary evolution of the Tertiary of Eastern Sabah, Northern Borneo. PhD Thesis, University of London, 456 p.
- Omang, S.A.K. & Barber, A. J., 1996. Origin and tectonic significance of the metamorphic rocks associated with the Darvel Bay Ophiolite, Sabah, Malaysia. In: Hall, R. & Blundell, D.J. (Eds.), Tectonic Evolution of SE Asia. Geological Society London Special Publication, 106, 263-279.
- Parkinson, C. D., Miyazaki, K., Wakita, K., Barber, A. J. & Carswell, D.A., 1998. An overview and tectonic synthesis of the pre-Tertiary very-high-pressure metamorphic and associated rocks of Java, Sulawesi and Kalimantan, Indonesia. Island Arc, 7, 184-200.
- Pieters, P.E., Trail, D.S. & Supriatna, S., 1987. Correlation of Early Tertiary rocks across Kalimantan. Proceedings Indonesian Petroleum Association 16th Annual Convention, 291-306.
- Rangin, C. & Silver, E.A., 1991. Neogene tectonic evolution of the Celebes Sulu basins: New insights from Leg 124 drilling. In: Silver, E.A., Rangin, C., von Breymann, M.T. *et al.* (Eds.), Proceedings of the Ocean Drilling Program, Scientific Results, 124, 51-63.
- Rangin, C., Bellon, H., Benard, F., Letouzey, J., Müller, C. & Tahir, S., 1990a. Neogene arc-continent collision in Sabah, N. Borneo (Malaysia). Tectonophysics, 183, 305-319.
- Rangin, C., Jolivet, L. & Pubellier, M., 1990b. A simple model for the tectonic evolution of southeast Asia and Indonesia region for the past 43 m.y. Bulletin de la Société géologique de France, 8, 889-905.
- Rangin, C., Le Pichon, X., Mazzotti, S., Pubellier, M., Chamot-Rooke, N., Aurelio, M., Walpersdorf, A. & Quebral, R., 1999a. Plate convergence measured by GPS across the Sundaland/ Philippine sea plate deformed boundary: the Philippines and eastern Indonesia. Geophysical Journal International, 139, 296-316.

- Rangin, C., Spakman, W., Pubellier, M. & Bijwaard, H., 1999b.
 Tomographic and geological constraints on subduction along the eastern Sundaland continental margin (South-East Asia).
 Bulletin de la Société Géologique de France, 170, 775-788.
- Reinhard, M. & Wenk, E., 1951. Geology of the colony of North Borneo. Bulletin of the Geological Survey Department of the British Territories in Borneo, 1, 160 p.
- Replumaz, A. & Tapponnier, P., 2003. Reconstruction of the deformed collision zone between India and Asia by backward motion of lithospheric blocks. Journal of Geophysical Research, 108, doi:10.1029/2001JB000661.
- Replumaz, A., Karason, H., van der Hilst, R.D., Besse, J. & Tapponnier, P., 2004. 4-D evolution of SE Asia's mantle from geological reconstructions and seismic tomography. Earth and Planetary Science Letters, 221, 103-115.
- Sandal, S.T. (Ed.), 1996. The Geology and Hydrocarbon Resources of Negara Brunei Darussalam, Syabas, Bandar Seri Begawan, Brunei Darussalam. 243 p.
- Sandwell, D.T. & Smith, W.H.F., 2009. Global marine gravity from retracked Geosat and ERS-1 altimetry: Ridge Segmentation versus spreading rate. Journal of Geophysical Research, 114, B01411, doi:10.1029/2008JB006008.
- Simons, W.J.F., Socquet, A., Vigny, C., Ambrosius, B.A.C., Abu, S.H., Promthong, C., Subarya, Sarsito, D.A., Matheussen, S., Morgan, P. & Spakman, W., 2007. A decade of GPS in Southeast Asia: Resolving Sundaland motion and boundaries. Journal of Geophysical Research, 112, B06420, doi:10.1029/2005JB003868.
- Suggate, S., 2011. Provenance of Neogene sandstones of Sabah, northern Borneo. PhD Thesis, Royal Holloway University of London, 441 p.
- Suggate, S.M. & Hall, R., 2013. Using detrital garnet compositions to determine provenance: a new compositional database and procedure. In: Scott, R.A., Smyth, H.R., Morton, A.C. & Richardson, N. (Eds.), Sediment Provenance Studies in Hydrocarbon Exploration and Production. Geological Society London Special Publication, 386, 373-393.
- Suggate, S.M., Cottam, M.A., Hall, R., Sevastjanova, I., Forster, M.A., White, L.T., Armstrong, R.A., Carter, A. & Mojares, E., 2014. South China continental margin signature for sandstones and granites from Palawan, Philippines. Gondwana Research, 26, 699-718.
- Tan, D.N.K., 1979. Lupar Valley, west Sarawak. Geological Survey of Malaysia Report, 13, 159 p.
- Tan, D.N.K., 1982. Lubok Antu Melange, Lupar Valley, West Sarawak: a Lower Tertiary subduction complex. Bulletin of the Geological Society of Malaysia, 15, 31-46.
- Tan, D.N.K. & Lamy, J.M., 1990. Tectonic evolution of the NW Sabah continental margin since the Late Eocene. Bulletin of the Geological Society of Malaysia, 27, 241-260.
- Tang, Q. & Zheng, C., 2013. Crust and upper mantle structure and its tectonic implications in the South China Sea regions. Journal of Asian Earth Sciences, 63, 510-525.
- Tapponnier, P., Peltzer, G. & Armijo, R., 1986. On the mechanism of collision between India and Asia. In: Coward, M.P. & Ries, A.C. (Eds.), Collision Tectonics. Geological Society of London Special Publication, 19, 115-157.
- Tate, R.B., 1991. Cross-border correlation of geological formations

in Sarawak and Kalimantan. Bulletin of the Geological Society of Malaysia, 28, 63-96.

- Tate, R.B., 2001. The geology of Borneo Island. Geological Society of Malaysia.
- Taylor, B. & Hayes, D.E., 1980. The tectonic evolution of the South China Sea Basin. In: Hayes, D.E. (Ed.), The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands. American Geophysical Union Geophysical Monograph Series, 23, 89-104.
- Taylor, B. & Hayes, D.E., 1983. Origin and history of the South China Sea Basin. In: Hayes, D.E. (Ed.), The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands, Part 2 American Geophysical Union Geophysical Monograph Series, 27, 23-56.
- Tongkul, F., 1991. Tectonic evolution of Sabah, Malaysia. Journal of Southeast Asian Earth Sciences, 6, 395-406.
- Tongkul, F., 1993. Tectonic control on the development of the Neogene basins in Sabah, East Malaysia. Bulletin of the Geological Society of Malaysia, 33, 95-103.
- Tongkul, F., 1994. The geology of northern Sabah, Malaysia: Its relationship to the opening of the South China Sea basin. Tectonophysics, 235, 131-137.
- Tongkul, F. & Chang, F. K., 2003. Structural geology of the Neogene Maliau Basin, Sabah. Bulletin of the Geological Society of Malaysia, 47, 51-61.
- van Bemmelen, R.W., 1949. The Geology of Indonesia. Government Printing Office, Nijhoff, The Hague. 732 p.
- van Hattum, M.W.A., 2005. Provenance of Cenozoic sedimentary rocks of northern Borneo. PhD Thesis, University of London, 467 p.
- van Hattum, M.W.A., Hall, R., Pickard, A.L. & Nichols, G.J., 2006. SE Asian sediments not from Asia: Provenance and geochronology of North Borneo sandstones. Geology, 34, 589–592.
- van Hattum, M.W.A., Hall, R., Pickard, A.L. & Nichols, G.J., 2013. Provenance and geochronology of Cenozoic sandstones of northern Borneo. Journal of Asian Earth Sciences, 76, 266-282.
- Wilford, G.E. & Kho, C.H., 1965. Penrissen area, west Sarawak, Malaysia. Malaysian Geological Survey Borneo Region, Report 2, 195 p.
- Williams, P.R., Johnston, C.R., Almond, R.A. & Simamora, W.H., 1988. Late Cretaceous to Early Tertiary structural elements of West Kalimantan. Tectonophysics, 148, 279-298.
- Wilson, R.A.M. & Wong, N.P.Y., 1964. The geology and mineral resources of the Labuan and Padas Valley area, Sabah, Malaysia. Geological Survey of Malaysia, Borneo Region, Memoir 17, 150 p.
- Wolfenden, E. B., 1960. The Geology and Mineral Resources of the Lower Rajang Valley and adjoining areas, Sarawak. British Territories Borneo Region Geological Survey Department, Memoir 11, 167 p.
- Wolfenden, E.B. & Haile, N.S., 1963. Semantan and Lundu area, West Sarawak. British Borneo Geological Survey Report, 1, 159 p.
- Wong, Y.L., 2011. Stratigraphy of the Ransi Member of the Middle Eocene to Oligocene Tatau Formation in the Tatau-Bintulu area, Sarawak, East Malaysia. MSc Thesis, University of Malaya, Kuala Lumpur. 256 p.

Manuscript received 9 June 2016 Revised manuscript received 26 August 2016 Manuscript accepted 24 November 2016